

INDEX NUMBER

Vol. II

DECEMBER, 1916

No. 12

The
**INTERNATIONAL
JOURNAL
OF
ORTHODONTIA**

*A Monthly Journal Devoted to the Ad-
vancement of the Science of Orthodontia*

MARTIN DEWEY, D.D.S., M.D., Kansas City, Mo.
Editor-in-Chief

H. C. POLLOCK, D.D.S., St. Louis, Mo.
Associate Editor

JAMES DAVID McCOY, D.D.S., Los Angeles, Calif.
Editor, Department of Dental and Oral Radiography

PUBLISHED BY
THE C. V. MOSBY COMPANY
SAINT LOUIS

\$3.00 Per Annum

Single Copies, 30 Cents

Copyright, 1916, by The C. V. Mosby Company.

LISTERINE

is an unirritating, non-caustic antiseptic solution, well adapted to many requirements in Dental Practice. It is the strongest and safest of the antiseptics available for daily use as a wash for the mouth and teeth.

LISTERINE

being slightly acid, is most valuable and effective in preventing the multiplication of various microorganisms which thrive in the normally alkaline saliva of the mouth.

LISTERINE

because of its slight acidity and aromatic flavor, stimulates the flow of saliva and thereby assists in maintaining a healthy equilibrium of the fluids of the oral cavity.

"The Dentist's Patient" is a booklet designed to impart useful information to the patient regarding the teeth and their care, and advocates frequent visits to the dentist's office. We publish and deliver these booklets free of cost in lots not exceeding 200 copies, imprinted with the dentist's name and address when desired.

LAMBERT PHARMACAL COMPANY
TWENTY-FIRST AND LOCUST STREETS, ST. LOUIS, MO.

The International Journal of Orthodontia

Editor: Martin Dewey, D.D.S., M.D.

VOL. II

ST. LOUIS, DECEMBER, 1916

No. 12

ORIGINAL ARTICLES

THE ROLE OF MOUTH INFECTIONS AND MOUTH ABNORMALITIES IN THE CAUSATION OF DISEASE*

BY EUGENE LYMAN FISK, M.D.

Director of Hygiene, Life Extension Institute, New York City.

THERE is no more important field of preventive medicine than that of the hygiene of the head. The more one studies the modern man—civilized, cultured, pampered, served by scientifically harnessed natural forces almost as was Aladdin by the slaves of the lamp—the more evident it becomes that disease, old age, and death fairly radiate from the cavities of the head.

Take, for example, the repeated acute infections having their origin in obstructed nasal passages lined with unhealthy mucous membranes poorly equipped to meet infection. Season after season the mass of the population suffer from these nasopharyngeal affections involving acute bacterial attack and resultant strain of organic tissue. In some cases death occurs from acute pneumonia, mastoiditis, brain abscess; in others chronic pulmonary disease or tuberculosis. Setting aside the effects of repeated frank, acute nasopharyngeal affections, rhinitis, grippe, tonsillitis, regarded as more or less normal experiences of the average individual, and also eliminating pneumonia and tuberculosis, we have to consider the effects of the establishment in the various cavities of chronic foci of infection. These cavities, tooth-sockets, accessory sinuses, tonsillar crypts, middle ear, etc., are, as Rosenau and others have shown, veritable culture tubes, where the streptococci and pneumococci go into training, as it were, before streaming out to attack those portions of the body where they can best thrive. Thus it is no mere figure of speech to say that in the head are the training camps for the most dangerous enemies of the human race. These microorganisms are among the chief causes of chronic disease and therefore of old age and death. As Mayo has expressed it, "Life is one long struggle with microorganisms."

*Read before the Eastern Association of Graduates of the Angle School of Orthodontia, May 5, 1916.

Deficiency diseases we undoubtedly have, like beriberi, scurvy, and probably pellagra, but I am firmly convinced that long before such stages of serious tissue impairment due to deficiency of vitamins or other necessary elements are reached, there are various degrees of deprivation that lessen the resistance of bacterial attack, substandard conditions of the body that pass unnoticed and unnamed until some organ obviously fails in its task, and then we call it disease.

Eliminating the so-called communicable diseases, such as typhoid and tuberculosis, we still have a tremendous death-rate from disease that is essentially due to infection. Chronic cardiorenal-vascular diseases are responsible for more than twenty-five per cent of the total death-rate. We know now that in all these organic affections, bacteria play a prominent and often the principal part. Injury, strains, poisons, endogenous or exogenous, all have their influence, but bacterial attack must be recognized at some stage of the game. I believe I am safe in saying that if the cavities in the head could be kept in a thoroughly healthy state, free from infection and their functions well maintained, at least four years could be added to the expectation of human life. This sounds like a little bit, but it means a tremendous reduction in the death-rate.

I would not minimize the role of the intestine as a great focus of subinfection, and of miscalled auto-intoxication. But it is not as guilty as formerly supposed. Indeed, in much of the damage that it does, it is merely an accomplice after the fact of focal head infection that has travelled downward.

The freedom with which bacteria range throughout the human organism has only lately been recognized. Subinfection, radiating from focal infection, must be recognized as one of the principal factors in organic impairment. Even in tuberculosis we find that the process of infection is not so simple as formerly supposed. The fact that tubercle bacilli can traverse the stomach and intestinal canal, penetrate the walls of the intestine without leaving any trace, and finally reach the bronchial glands or lungs, is an illustration of the losing fight that mankind is maintaining with the various forms of bacteria.

Since death comes to all ultimately, always in the form of a pathological state, I am justified in terming this a losing fight. What I have said regarding the hygiene of the head and the importance of the head cavities as potential factors in organic disease, emphasizes the tremendous responsibility upon those who are specializing in that region. It is the region par excellence for the entry of infection, as well as for nutritive material and oxygen.

It would be difficult to find an individual with absolutely normal and perfectly functioning head cavities. In using this expression, I am of course excluding the brain. Harry Campbell has recently stated that there are upwards of two hundred millions of carious teeth in the British Isles, and even a greater number of infected tooth sockets, while as to jaws, it is rare to meet a normal one. There is no reason to believe that these conditions are any better in this country. False teeth, long teeth, irregular teeth, infected gums, asymmetric jaws, are the rule rather than the exception. The field, therefore, is an almost limitless one, and the chief responsibility upon the specialist, as upon all others who are working in modern medicine, is to begin as early as possible in the life of the individual to detect abnormalities of these cavities, correct them, and protect the individual from future impairment.

The protection of the deciduous teeth falls to the parent and the dentist, as a part of mouth hygiene. While success in this direction will limit the need for the orthodontist, there will always be plenty for him to do in correcting those forms of irregularity and asymmetry due to heredity or other causes. Also, on the one hand, the orthodontist can lessen the need for rhinological work and septal operations, while, on the other hand, the rhinologist can in his own field assist the individual to a healthy and more vigorous development of the upper jaw and lessen the need for orthodontia in that region, also the general practitioner or hygienist can, by regulating the diet and training the child to eat properly, prevent dental irregularities.

It is desirable to carry to the profession at large the knowledge of the possibilities of orthodontia, knowledge that it performs far more than a cosmetic service, that while the relief of deformities has a very high value from a social and even health standpoint, these tooth irregularities contribute very largely to mouth infection and hence to organic disease. It seems true that the modern mouth is degenerate in its parts.

There may be members present who have studied this question more deeply than I have, in which case I should be more than glad to have additional evidence on this point. There is, however, abundant evidence to show that in a state of nature man's jaws are more vigorous and symmetrically developed, and that he is also comparatively free from other head abnormalities and asymmetries so common among the civilized. Whether early mental development and educational strain have anything to do with this matter, as some authorities have suggested, I am unable to say; but heredity, and deficiencies in the diet in infancy, as well as the long list of hygienic errors that characterize civilized life, must be given due weight.

These possibilities of head infection emphasize what is true of all the other regions of the body: namely, that from infancy on, they should be regularly inspected for abnormal conditions. I differ with Dr. Cabot, who claims that these inspections are of limited value unless made by specialists in each region. That is not feasible as applied to the mass of the public, but under a proper routine, as advocated by the Life Extension Institute, a well-trained general practitioner can make such periodic examinations to detect the need for further special attention.

We must sharply distinguish between the clinic for established disease and its treatment, and this sifting process by which the early signs of disease are noted and the patient shunted to whatever special clinic or clinician his needs may require.

It is amazing what an automatic, systematic sieve of this description will reveal. Indeed, it will bring out more evidences of impairment than is usually the case when a man drifts aimlessly around from one specialist's office to another. The examiner who is trained and educated to make such a survey performs the function of a sort of medical air-man, who searches out the enemy's weaknesses and gives information by which successful attack can be made.

The following figures from the examinations of the Institute will give you an idea of the operation of this principle:

RESULTS OF PHYSICAL EXAMINATIONS OF INDUSTRIAL AND COMMERCIAL WORKERS
BY THE LIFE EXTENSION INSTITUTE COMPARED.

	Industrial.	Commercial.
	Average age	Average age
	31.6 years.	27.8 years.
Free from physical defects or unhealthful living habits	0.46%	0.81%
Slight impairments or errors in living habits	27.16%	22.96%
Moderate impairments, needing medical, dental, or hygienic treatment	67.15%	71.81%
Important impairments, requiring close medical supervision and treatment	5.20%	4.28%

It will interest you to know that 69 per cent of these people examined showed need of dental attention, and that among the number showing that need, organic impairment was found to the extent of 22 per cent in excess of those without dental impairment.

I think you will agree with me that this specialized assistance, such as workers in the field of orthodontia are able to render, should not be left to the haphazard initiative of patients. The detection of these abnormalities that require your aid should be a systematic, routine affair, applied to the whole population. It is then up to you to provide facilities by which the benefits of this treatment shall not be limited to the idle rich, but extended by some means to all who need it.

INTRAMAXILLARY ANCHORAGE

BY MARTIN DEWEY, D.D.S., M.D., KANSAS CITY, MO.

*Professor of Dental Anatomy and Orthodontia, Kansas City Dental College;
President of The Dewey School of Orthodontia.*

I NTRAMAXILLARY anchorage may be defined as that type of anchorage in which the resistance necessary to overcome a malposed tooth is obtained from a tooth located in the same arch.

Intramaxillary anchorage is divided according to two plans. The first plan is based upon the number of teeth employed in the resistance, and the second upon the manner in which the resistance is obtained. We speak of intramaxillary anchorage as being simple or compound, according to the number of teeth employed, the latter being sometimes called "reinforced." In the seventh edition of Angle's book on "Malocclusion of the Teeth," and in my work, "Practical Orthodontia," the word "simple" was used to designate anchorage according to method and is defined as being "that form of anchorage where the resistance is obtained from a large tooth more favorably located." Owing to the great amount of confusion that has arisen in the use of the term "simple," it was decided to change the term so as to refer to numbers instead of plans. Therefore, in reading this article it will be well to remember that the terms have been changed in that respect; and that "simple anchorage" is here used as referring to the number of teeth from which resistance is obtained and not to the size and location of the

anchor teeth. This may cause more or less confusion in the minds of some, but when using the term "simple" in regard to method and "compound" in regard to numbers, the confusion continually exists in the minds of students, because in the general use of these two words they are opposite in meaning.

The term "simple anchorage" is used in this paper as referring to numbers and that form of anchorage obtaining resistance from a single tooth. In Fig. 1, taken from the seventh edition of Angle's book, we have a very good illustration of a case of simple intramaxillary anchorage in which a second deciduous molar is used to overcome the resistance necessary to move the incisor. No provision has been made in the construction of the appliance to enlist




Fig. 1.




Fig. 2.

the support of any of the other teeth, however, it must be remembered that, although the appliance is attached only to one tooth, owing to the intimate connection of the fibers of the periodontal membrane, there will necessarily be some force transmitted or taken up by the surrounding tissue and proximating teeth.

Compound or reinforced intramaxillary anchorage is that form of anchorage where two or more teeth are used to overcome the resistance of the malposed tooth. The construction of this appliance provides for the embracing of two or more teeth. An example of intramaxillary anchorage is shown in Fig. 2 which includes the use of the jackscrew, with attachments made to the incisor which supports the premolar carrying the anchor band; and by the proximal contact,

all of the incisors, to a certain extent, become anchor teeth. This compound or reinforced anchorage may be produced by banding more than one tooth, or it may be produced by constructing or applying an auxiliary appliance attached to a spur soldered to the band so as to embrace or rest against more teeth than would be influenced by the band itself. Fig. 3 shows a type of reinforced or compound primary intramaxillary anchorage in which a spur has been soldered to the lingual surface of the anchor band on the second premolar, the spur resting against the first premolar and molar with the result that the force exerted on




Fig. 3.




Fig. 4.

the premolar as an anchor tooth is also exerted upon the proximating molar and premolar. This presents a type of reinforced or compound intramaxillary anchorage in which the same appliance is used as shown in Fig. 4, which represents a case of simple intramaxillary anchorage. In considering simple and compound intramaxillary anchorage, it must be remembered that it makes no difference which teeth are employed, how the appliance is attached, or what kind of an appliance is used; the classification being based solely upon the number of teeth mechanically attached to the appliance. This classification, simple and compound, must not be confused with the method of attachment which brings us to another classification of anchorage. A great many more examples of simple

and compound intramaxillary anchorage might be shown, but the above are sufficient for the purpose of illustrating these two types of anchorage.

METHOD OF ANCHORAGE.

In taking up the consideration of methods of anchorage, we have to deal with the plans of obtaining the resistance to overcome the malposed teeth. We have three plans in intramaxillary anchorage which are known as primary, stationary, and reciprocal. In the classification of anchorage as given in my "Practical Orthodontia," the term used is "primary anchorage" when referring to numbers because the term "simple anchorage" has already been appropriated by other writers with reference to method. As mentioned in the first part of this paper, owing to the confusion that has arisen in the minds of students, it has been decided that it would be better for orthodontia to use the word "primary" in reference to method and let it describe the same type of anchorage as Angle has described under "simple" anchorage.

Simple primary intramaxillary anchorage is then defined as that type of anchorage in which the resistance necessary to overcome the malposed tooth is derived from a larger tooth or one more favorably located. An example of simple primary intramaxillary anchorage is shown in Fig. 4.

Compound primary intramaxillary anchorage is that form in which the force necessary to produce movement of the malposed tooth or teeth is overcome by two or more teeth larger or more favorably located. In compound primary intramaxillary anchorage, if it is necessary to move a tooth which requires a large amount of force, we obtain that force by utilizing two or more teeth and thus get the resistance from an increased number and a more favorable location. In compound primary intramaxillary anchorage, it makes no difference whether or not more than one tooth carries a band, only so far as we enlist more than one tooth as the anchor tooth. Primary anchorage has also been described as that type in which, if sufficient force is brought to bear upon the anchor tooth, it will tip through the process instead of moving bodily. The primary intramaxillary anchorage defined by Angle under the term "simple anchorage" is that "form in which, first, resistance is overcome by the anchor tooth or teeth of larger size or of a more favorable location, and second, the form of attachment to both anchor tooth, and teeth to be moved, is hinged or pivoted, admitting of tipping of both in their sockets." In this definition he includes both simple and compound intramaxillary anchorage and discusses the point that makes intramaxillary anchorage of the primary type different from stationary anchorage. Primary intramaxillary anchorage is the easiest type of anchorage obtained because, regardless of how the appliance is attached to the anchor tooth when we depend only upon the resistance of the anchor tooth, we necessarily must have primary intramaxillary anchorage. However, with this type of anchorage it must be remembered that no extra resistance is obtained by the construction or the attachment of the appliance; if sufficient force is brought to bear upon the anchor tooth it will necessarily be tipped.

With primary anchorage, the knowledge of the alveolar process and the

anatomical surroundings must be employed in order to obtain as favorable a location of the anchor tooth as possible. The relative sizes of different teeth must also be considered as well as the different degrees of ease with which the teeth move in different directions. It must be remembered that a molar can be moved mesially with greater ease than it can be moved distally. Upper molars as a rule, can be moved buccally more easily than they can be moved lingually. The reverse is true of the lower molars, especially the second and third molars. The upper incisors as a rule, will move labially more easily than they will move lingually. When speaking of the different resistance offered by different teeth when moving in different directions, it must be remembered that we are considering conditions in a normal alveolar process and with normal attachment to the teeth. One of the problems of anchorage which we encounter is the fact that in certain individuals the alveolar process is not normal as the result of which teeth move with much less force than they will in other mouths. An example of faulty alveolar process is found in those patients that have rickets or possess a rickety diathesis. In those cases, it is almost impossible to tell with any degree of certainty which teeth will move as a result of the application of force. For instance, if the alveolar process around the malposed tooth, or the one which is to be moved, is more normal than that around the anchor tooth, the anchor tooth will be displaced before the malposed tooth will move. Therefore, a set rule laid down for the movement of anchor teeth or malposed teeth as a result of anatomical resistances, can be applied only in individuals who have normal alveolar process. Owing to the fact that primary anchorage depends only upon the position of the anchor tooth and is not dependent upon the construction of the appliance so as to increase the anchorage, and owing to the fact that in a large number of cases the resistance of the tooth is not sufficient to produce tooth movement, it is necessary to evolve another type of intramaxillary anchorage which is described as "stationary anchorage."

Stationary anchorage as described by Angle is "that form in which the attachment to the anchor tooth is essentially rigid, so that tipping is impossible, and if it is moved at all, it must be dragged bodily through the alveolar process in the upright." This form of anchorage, according to Angle's writing, was first introduced in the second edition of his work, and he also refers to it in *Items of Interest*, December, 1887. He gives credit to Barrett "who first made use of this form of anchorage, though in an imperfect manner, by means of a vulcanite plate entirely covering the vault of the arch and molars, but this, of course, did not admit a strict rigidity of attachment." Stationary anchorage, however, was evolved by the early writers because of the difficulties encountered in producing certain tooth movement when depending entirely upon the "size and location of the anchor tooth." Owing to the fact that stationary anchorage differs from primary anchorage in the form of attachment, a better definition than the one given, is—stationary anchorage is that form in which the appliance is so constructed and attached to the anchor tooth that if the anchor tooth moves at all, it will be dragged bodily through the process. There are two features involved in the stationary anchorage which are not employed in primary anchorage; namely, the appliance must be constructed rigidly and must be rigidly attached to

the anchor tooth. This appliance described in Angle's work shows the use of the screw similar to the one shown in Fig. 5. Now, in order to have stationary anchorage with the traction screw, the traction screw must be so rigidly attached, and so rigidly constructed, that there will be no bending in that portion of the attachment on the canine and on the molar. It must also be so accurately adjusted that there will be no play between the tube that is soldered to the molar




Fig. 5.

band and the portion of the screw that extends to the canines. In order to insure the rigidity in the use of the stationary anchorage, there should be as much of a point of bearing between the fixed portion of the appliance and the moving portion of the appliance as is possible. In other words, all tubes which are rigidly attached to the bands on the anchor teeth should be as long as possible to receive the portion of the appliance that is going to be the active part.

In the next installment of this paper we will take up the various methods of obtaining stationary anchorage with the alignment wire.

A "PATENT" SUGGESTION

BY HARRY P. BEASER, FRESNO, CALIF.

SINCE the time of the invention of the porcelain crown, the subject of patents has occupied the attention of the dental profession to a great extent.

From the beginning of time men have progressed by invention, which is nothing more nor less than the discovery of a new idea, and laws have been made with the object of protecting the inventor so that he can reap the benefits therefrom.

There are, however, a great number of men whose only desire is to create something new and useful, and the big majority of men in the dental profession belong, I think, to this class. But owing to the fact that there are those who are unscrupulous enough to take advantage of this generosity, these men have been compelled to protect their inventions by law when they had no wish to get any financial gain therefrom.

I would therefore suggest that the National Dental Association take the matter of patents into their hands and either create a board commission to look after this subject, or make it a part of the Research Commission. If this were done, men in all branches of the profession, and the public at large, could send their inventions to this department of the National Dental Association knowing that no individual would get the benefit of the idea for his personal gain to the exclusion of all others.

This patent board could look into the merits of the invention, and if found worthy of consideration, could obtain a patent in the name of the National Dental Association, giving the originator due credit on their records and in the *Journal of the National Dental Association*.

The dental profession, especially in America, has reached its high standard of efficiency through the inventive genius of its members; and there have been many new ideas of great value which have never gone beyond the door of the office because the founders did not wish to undergo the expense of obtaining a patent, or take the chance of losing them by giving them to the profession, and then have some one else get the patent on their article and make them pay royalty for the creation of their own minds; also for the reason that they did not wish to place themselves in the position of trying to collect a royalty from other members of the profession.

If this suggested system had been a fact a number of years ago, there is no doubt that a great deal of trouble would have been avoided; e. g., previous to the time of the discovery of the method of making the cast gold inlay, the disappearing wax models had been used for other purposes. If this had been recorded and taken care of by the National Dental Association at the time of its real discovery, no process patents would ever have been given and no trials for infringements would ever have entered the courts.

This may all be an old idea and I do not claim originality for it, but from investigations I have been able to make I have not discovered it. I believe, however, that it could be worked out to advantage even if it cost the profession a few cents per head more than their present dues, for all would reap the benefits alike and many disagreeable and costly law proceedings would be eliminated. It would certainly be a cure for a lot of unprofessionalism and trend toward bringing all men closer together for the common good.

TREATMENT OF CERTAIN TYPES OF MALOCCLUSION WITH THE AID OF BRIDGE-WORK

BY H. E. BLILER, D.D.S., CHICAGO, ILL.

THE correction of malocclusion must necessarily be considered from different angles and there are certain factors entering into the treatment of one case which may not influence another—the time required to correct the malocclusion, the inflammation created, the inconvenience to patients, their financial status, and the best mode of procedure to obtain quick and lasting results. Every general practitioner should be able to correct simple cases, especially where it is not possible to refer the patient to a specialist, as in the smaller towns and rural districts. It is a well known fact that cases can be treated successfully at a certain age while at a more advanced period of life the treatment of them is much more



Fig. 1.—Showing models of case before and after treatment. Fifteen days was allowed for shrinkage or absorption of the V-shaped arch.




Fig. 2.—Gold crown was placed on six-year molar.

troublesome even if not prohibitive. It must also be remembered that certain people can give a considerable length of time to the treatment of malocclusion while others have a limited amount of time and cannot devote several years to the treatment, if there is any other means of accomplishing an improvement in a shorter space of time. It must also be remembered that in some instances the financial inability of the patient to pay for a long treatment, even though the result meant the ideal condition, often causes the operator to decide in favor of a plan whereby he accomplishes results in much less time even though the results are not as ideal as those that require a greater amount of time.

Fig. 1 shows the model of a case of a girl fifteen years of age who was not able to spend the necessary time and money to have the case rectified by orthodontic methods. Also the amount of deformity which is shown in the teeth would have necessitated some bridge-work in the end. Fig. 1 was made from an impression taken by having the patient bite on wax. It shows that the lateral incisors




Fig. 3A.—Gold crown was placed on six-year molar.



Fig. 3B.—Patient was a mouth-breather, not able to keep her lips closed on account of prominent centrals.



Fig. 4.



Fig. 5.

Figs. 4 and 5.—Gratifying results obtained by mechanical skill, when nature helps.

were small, consequently something was required in order to improve the lateral incisors as well as to correct the protrusion of the centrals. Both centrals, which were oversized, were extracted and the pulp was killed in the two undersized lateral teeth. February 29th I cut off the lateral teeth and made a four-tooth bridge, using porcelain interchangeable teeth, with the half-band attachment,

finishing March 3rd, giving the appearance as shown in Fig. 2. One of the central incisors was broken off, the lateral incisors were undersized, and it necessarily made a very unpleasing type of malocclusion, which is shown in Fig. 3. Fig. 4 shows the appearance of the patient's face and lips in repose after the case was treated, while Fig. 5 shows the appearance of the patient's face while smiling. In my opinion it was the correct solution of the case regardless of her ability to pay more or less.

We are aware of the fact that the extraction of an anterior tooth in the correction of malocclusion is to be avoided, but here was a case where the centrals were deformed by being oversized, one of them was broken off, the lateral incisors were small, even if the strictly orthodox plan of orthodontic treatment was followed bridge-work and restoration would have been necessary. This meant time and money as a decided factor, consequently, the method of extracting the two large incisors, and adjusting a four-tooth bridge by using the small lateral incisors as abutment, was decided upon. In fact I am very much in doubt if a better appearance could have been accomplished considering the patient we had to work on, and the condition in the beginning, if a different plan had been followed. However, in the use and the construction of the bridges for the correction of malocclusion, they must be constructed according to anatomical laws, the force of the occlusion must be observed, the operator must be positive that the constructed bridge will be better in appearance than the deformed and mutilated teeth which are extracted.

I get good results many times by extracting the lower central in badly crowded conditions rather than torturing the patient by trying to widen the arch, or by extracting the first bicuspid and trying to move several teeth over. By extracting the central, the teeth will straighten themselves out, and it is good practice for poor patients, who otherwise would have crowded teeth all their lives. I extracted a lower lateral tooth for a man, 37 years old, and the teeth straightened, without any appliances, in a year's time.

BANDS OR CROWNS FOR TEMPORARY MOLARS

BY W. A. COSTON, D.D.S., TOPEKA, KANSAS.

THE difficulties—discomfort and sometimes pain attending the proper fitting and adjusting of anchor bands on molars, especially the temporary molars, is sometimes quite annoying to both the little patient and the dentist, and the fact that they sometimes do not stay on—that they loosen, come off, or are driven into the soft tissue by the force of occlusion, and cause inflammation of the gingivæ, if not the peridental membrane, or sometimes cause etching of the enamel, is also quite annoying. All of these things happen to the "busy dentist," and the following method, which has proved very satisfactory to the author, is suggested for the relief and the comfort of the little patient.

The indirect method of making various restorations has been a great benefaction to dentists—especially the general practitioner, and a great factor in all

prosthesis—as we have learned that crowns, inlays, clasps, and almost every appliance or restoration that is to be cemented to place, can be made with more accuracy, generally speaking, and certainly with more facility than we have been able to do by any other method. This idea applied to the making and fitting of anchor bands is quite as practical as for the making of bands or crowns for



Fig. 1.—Thin copper band trimmed to festoon of gum.



Fig. 2.—Impression of tooth in compound.



Fig. 3.—Impression with celluloid strip wrapped around it.




Fig. 4.—Impression invested in plaster.



Fig. 5.—Copper amalgam model of tooth.



Fig. 6.—Band soldered with lap joint.



Fig. 7.—Band swedged.



Fig. 8.—Cusp swedged for band.



Fig. 9.—Band with cusp soldered and cut out for occlusion.



Fig. 10.—Crown with tube and lug.

any other purpose. After the necessary separation for the application of a band, an impression of the individual tooth is taken, using a thin loosely fitting copper band, trimmed to the festoon of the gum (Fig. 1), and made thin enough to pass into the approximal space without crowding, which is filled with soft modelling compound, and forced to place on the tooth. When chilled it is removed

(Fig. 2), encompassed with a celluloid strip (Fig. 3), and sunk into thin plaster (Fig. 4.) When the plaster is set the impression is filled with copper amalgam, mixed to a soft consistency as in indirect work. When the amalgam has set, the plaster containing the model is slightly warmed (for safety), broken open, and the amalgam model cleaned, using chloroform to remove any traces of modelling compound.

We now have a practical duplicate of the crown of the tooth in amalgam (Fig. 5) upon which a band of any description may be fitted with facility. In the method suggested, band material of a suitable gauge is cut from a tin foil pattern, and soldered, with lap joint (Fig. 6), and fitted to the model (joint on opposite side from the tube or other attachment), trimmed cusp-high or sometimes overlapping the cusps and marginal ridges a little. If necessary the band is slit a little and lightly burnished over and into the developmental grooves and margins to prevent displacement, and is then swedged to place in cylindrical swedge with vulcanite rubber. The band is then removed and the cusps trimmed high, allowing the band to lap over the occlusal surface at any place where it will not conflict with occlusion as in the buccal and lingual developmental groove (Fig. 7). This can now be reinforced with solder and used as a plain band, if desired, or, a thin piece of metal can be cut and annealed and swedged right over the band and model which will be forced down into the fissures and sulci and over the edge of the band (Fig. 8); then remove and solder to the band and finish. The crown, for such it is, will slip into place on the tooth, many times with a spring, and will interfere very slightly with occlusion, which interference can be remedied by grinding or cutting away the interfering metal from the occlusal surface, leaving enough overlapping the occlusal surface to prevent the band being forced gingivally into the tissue, should it loosen (Fig. 9).

The tubes or attachments can be soldered (Fig. 10), the band electroplated, and cemented to place with thin well-mixed cement, which causes it to stay on, that is, reasonably well, without any chance of gingival inflammation, if the cement is all cleaned off as it should be. The band can be any gauge desired for strength, but 34-gauge is surely thick enough. The cusp can be much thinner and in cementing the pressure should be on the band—or attachment thereto, and even a little lug can be soldered to the lingual for the purpose, as shown on Figs. 9 and 10, but pressure should not be on the soft and thin cusp.

If in the opinion of the operator the band swedged directly in the model fits too closely, first swedge and burnish tin foil, No. 60, over the model and then proceed as before mentioned—the result is accurate but loosely fitting.

A REMOVABLE RETAINING APPLIANCE

BY HARRY P. BEASER, D.D.S., FRESNO, CALIF.

RETAINING appliances are mechanical devices placed upon the teeth which have been moved from positions of malocclusion to hold them in a straight line of occlusion until the natural forces have been established which will maintain them in their proper position. Retaining appliances are divided into the fixed and removable, each one of which possesses advantages and disadvantages. The fixed appliance can probably be made a little more delicate than the removable appliance, but it holds the teeth more rigidly, as a rule, and does not admit of being removed so that the teeth can receive prophylactic care during retention. In the mouths of some patients who have always kept their teeth clean and whose teeth are not susceptible to cavities, a fixed appliance may be used without any ill results. However, in those mouths where prophylactic care must be given the teeth, an appliance which is removable, so that the parts can be kept clean, is better, and will be found a great advantage.

The appliance which is shown in the cuts is a retaining appliance constructed so that it will fill nearly all of the requirements of practically any case of malocclusion. It can be made equally adaptable to a mesiocclusion, neutroclusion, or a distocclusion case. The accompanying cuts show it constructed for a distocclusion case. One of the principal advantages it has over other mechanical retainers is that it is constructed as separate parts so that any one may be removed at any time without interfering with the rest. If a band becomes loosened on any tooth, it can be recemented without removing the entire appliance or running the risk of changing the shape. Each band is made and attached separately the same as an appliance constructed for treatment, thus reducing the difficulties to a minimum and at the same time producing the maximum of efficiency. It is also a well known fact that single bands are more easily set and more accurately fitted to the teeth than are three or four bands which are attached to one appliance, which necessitates the cementing of all at the same time. The appliance may be made either active or passive at any time or modified or added to at the will of the operator with the very least amount of time and trouble. Taken altogether this will come near to occupying that position so much required by orthodontists, that is, a retaining appliance which can be made into an active regulating appliance, whenever the emergency arises, and all orthodontists know this emergency arises very often. The maintenance of tooth position has caused orthodontists more trouble than all other intricacies in science. This fact is shown by early investigators who wrote at length describing all manners of appliances for the movement of the teeth with very little or nothing about retention. It is plain to admit that there was very little understood about the physiological changes which took place during the treatment of malocclusion.

It is very thoroughly understood at the present time that a correct diagnosis is absolutely necessary, not only to outline the course of treatment, but the retention as well, yet at the present time there is less difficulty experienced in

treatment, than in maintaining the teeth in the line of occlusion after they are placed there. This fact I believe is due to the construction of too rigid appliances for retention.

Nature if given a crutch will invariably lean upon it, but if forced to stand, will build a foundation for herself. This fact is proved in hundreds of




Fig. 1.




Fig. 2.

ways, for instance, the horse-riding Indians of the plains almost lost the art of walking. The body of a man who sits continuously, conforms itself to the shape of the chair, and the muscles all but lose their normal functions. A strong healthy person may start the use of drugs to stimulate some part of the system and in a short time the system requires the drug in order to perform its duty.

Thus it is with the teeth, and if a too rigid retainer is used, the teeth will lean upon it, and the tissues will atrophy instead of building strength. Therefore that appliance which will permit a free movement of a tooth in any direction, only acting as a guide to its position in the dental arch and permitting all of the forces of occlusion to act in harmony, is the one which will produce the best results.

The cuts here show the appliance fitted to a case of bilateral distoclusion which has been produced by mouth breathing. The materials used are the same as for any other appliance. Plain bands are made for the molars and canines,




Fig. 3.




Fig. 4.

upper (Fig. 1), and lower (Fig. 2). A 16 gauge vertical tube is soldered to the lingual side of the molar bands and a slot cut on the mesial surface about two-thirds of the length of the tubes from the occlusal end, large enough to permit a 20 gauge wire to slip in easily. When the wire is adjusted to the lingual surface of the teeth the ends are cut even with the inner side of the slot. The wire is then taken out and a piece of 16 gauge wire soldered perpendicular to each end making a T shape on each end of the 20 gauge wire. It is then slipped into the tubes and cut off smoothly. The T-shaped end on the 20 gauge wire which fits in the vertical tube prevents the molar from rotating.

In soldering the tubes to the molar bands they should be set at a slight angle so as to prevent the lingual wire from slipping out.

Small hooks are soldered to the linguo-cervical part of the canine bands (as in many other appliances) to engage the lingual wire and prevent it from slipping up. (Figs. 3 and 4.)

To antagonize the outward tendency of the upper anterior teeth a piece of




Fig. 5.




Fig. 6.

18 gauge tubing is soldered to the labial surface of the upper canine bands and a slot cut downward mesially and distally from the cervical end about two-thirds of the length, large enough to permit the easy access of a piece of 20 gauge wire. This can best be accomplished by filing the tubes about half through for the desired length and then soldering to the bands. A piece of 20 gauge wire will then be cut long enough to make a hook on each end and after being adjusted to the surfaces of the incisors will be slipped into place. (Fig. 5.) To prevent the rotation of the canine a perpendicular spur can be soldered to the labial wire to engage the tube on the canine band.

Intermaxillary rubbers will then be adjusted from the end of this labial wire to hooks on the lower molar band. (Fig. 8.) The ends of the tubes may be pinched together to prevent the wire from slipping out or being taken out by the patient, thus possessing all the qualities of the rigid soldered appliance without having any of its bad features.

If more freedom of movement is desired in the teeth to which the appliance is attached, the slots may be widened laterally at the points where the wires enter the vertical tube and the short 16 gauge pieces may be tapered at the ends giving the effect of a ball and socket joint.

Some writers have spoken disparagingly of the removable retaining appliances which various orthodontists have constructed from time to time, claiming that they were invented because of the inability of the operator to do free hand soldering with sufficient accuracy. This, however, I do not think worthy of much consideration, for a thing is not necessarily good because it is hard to make, or another bad because it is more simple; simplicity, efficiency, and durability being the most desired qualities that any appliance can possess.

Different sizes and kinds of material may be used as the operator sees fit to apply to the case in hand and modifications may be made to suit any class of work.

A REPORT OF AN ACCIDENT

BY H. B. HAMILTON, D.D.S., ITHACA, N. Y.

DENTAL accidents, more or less serious, are a frequent occurrence among children, and the following case is of interest because of the age of the child and the probable consequences on the growth of the mandible.

Betty H., aged eighteen months, knocked out three lower incisors in a fall down a flight of steps. The right central came out clean. The left central and lateral and a portion of the outer plate of process remained attached to the gum only, which, joining distal to lateral, was torn to the apex of the root. These two teeth lay in a horizontal position, the gum acting as a hinge and flopping back and forth with every movement of the tongue and lip.

At first, the case seemed hopeless, but it was found that the two teeth, when placed back in position, required but little to hold them. The canines were, of course, unerupted, and the age of the patient made it difficult to fit bands; so complete reliance for support was placed on the right lateral which was uninjured. A band was fitted to the lateral with a spur soldered to it sufficiently long to extend beyond the left lateral and cemented to place with the spur pressing lightly against the loosened teeth. Not quite enough pressure was obtained at first to hold the teeth back far enough, but this was corrected a few days later. Healing took place rapidly, and with the exception of recementing the piece a couple of times and removing three or four particles of process, no other treatment was required. A liquid diet was used for about four weeks. No attempt was made to replace the right central, and when last seen, about six weeks after the accident, the two teeth were in about the center of the space formerly occupied by the three teeth, and the canines just showing through the gum.

DEPARTMENT OF DENTAL AND ORAL RADIOGRAPHY

JAMES DAVID MCCOY, D.D.S., EDITOR,
LOS ANGELES, CALIF.

ELECTIVE LOCALIZATION OF THE STREPTOCOCCUS FROM A CASE OF PULPITIS, DENTAL NEURITIS AND MYOSITIS*

BY EDWARD C. ROSENOW, M.D., ROCHESTER, MINN.
The Mayo Foundation, Rochester, Minnesota.

LOCALIZED infections in or about the roots of teeth have already been considered in causal relationship to systemic disease and to neuralgia of the face. Experimental proof of the nature of this relationship, however, is still undetermined. In this paper I wish to record the history of a case in point and the results of experiments which appear to throw light on the subject.

Case 567.—Mrs. F. H. P., aged thirty-five years, had been subject to severe migraine for many years. For the past five or six years she has suffered from recurring attacks of neuralgia of the face, which began with severe pain and distinct swelling over the left upper jaw, spreading to the opposite side, with soreness in the teeth, especially in the upper jaw, and followed by intense pain in the left side of the head, neck and shoulders. During the last two or three years the attacks have occurred oftener and had grown so severe as to necessitate the frequent use of hypodermics of morphine; the attacks have ended with spasm of the muscles, and tenderness and swelling of the neck on the left side. Her tonsils were removed four years ago, but this did not relieve the condition. One year ago the second left upper molar showing a blind abscess at the root was extracted, the left maxillary sinus drained and a piece of the left turbinate removed, but without relief. She became extremely nervous, at times hysterical during the paroxysmal pain, and has had one or two spells of mental confusion suggesting *petit mal*. Previous to tonsillectomy, she had had for years one or more attacks of tonsillitis followed by rheumatic pains during the winter months.

Examination revealed a poor vasomotor tone, moderately firm muscles, fair nutrition, and the general appearance of a nervous woman. On January 13, the hemoglobin was 85 per cent; the leukocytes 9800. A Wassermann test of the blood proved negative. On February 17, the hemoglobin was 80 per cent; the erythrocytes 4,210,000; the leucocytes 5400. There were no signs of organic disease of the central nervous system. The examination of the

*Read before the American Association of Immunologists, Washington, D. C., May 11, 1916. Reprinted by special permission from *The Journal of Immunology*, Vol. I, No. 4, August, 1916. Received for publication June 4, 1916.

heart, lungs, abdomen, reflexes, urine, Roentgen examination of the jaws, roots of teeth, and blood pressure, were all negative. There was tenderness over the left infraorbital foramen and mental foramen, but no superficial tenderness of the overlying skin. There were tender nodules, which appeared to be enlarged glands in the posterior triangle of the left side of the neck. There was a clean tonsillectomy scar, a normal condition of the nose except moderate hypertrophic rhinitis, and a normal condition of the gums and teeth, except a tender dead first upper left molar which had been crowned, but from which the crown had been removed on account of the irritation of the gums. The maxillary sinuses were clear. Owing to the fact that each attack began with swelling of the left upper jaw opposite the dead tooth and in the region of the left infraorbital foramen, the tooth was extracted January 18. There was found a semi-lunar eroded area 2 mm. in diameter near the apex of the largest and inner fang. The surface of this was sterilized with a searing blade, and the root-canal drilled into from the apex with a dental burr. The pulp-cavity was filled with a foul-smelling pus. The erosion was situated so that it was impossible to be shown by the Roentgen ray. The pulp-cavity of the other two fangs was filled and obliterated and there were no erosions at the apices. There was a large cement filling directly opposite the dead pulp. Smears from the pus from the root showed a few Gram positive diplococci, diphtheroid bacilli and a few Gram negative bacilli. The primary cultures in ascites-dextrose agar and broth gave a pure culture of a slightly hemolyzing streptococcus. The anaerobic cultures on blood agar slants and in tall columns of ascites fluid containing sterile tissue, covered with paraffin oil, had a foul odor and many short chains of streptococci and numerous bacilli resembling *Bacillus fusiformis*.

Immediately after the extraction of the tooth, the patient developed an unusually severe attack but she gradually recovered after several exacerbations. On February 17, after having been free from pain for ten days and gaining rapidly in weight and strength, there was found a definite tenderness of the muscles in the posterior triangle on the left side of the neck and two distinct tender nodules just behind the posterior margin of the sternomastoid muscle. One of these tender nodules, thought to be a lymph-gland, and a portion of the deeper layers of the muscle, was excised. One-half of these tissues were immediately emulsified for cultures and the other half fixed in formalin-Zenker for sections. The excision of the fascia and muscle precipitated another violent attack of pain and spasm of the muscles of the left side of the neck. The cultures in tall columns of ascites plain broth of the emulsion of muscles showed a pure culture of a short chained streptococcus and those from the thickened fascia, streptococci and staphylococci.

Blood agar plates from the culture in ascites plain broth from the tooth pulp and from the muscle (injected into animals) showed pure cultures of streptococci producing a narrow hazy zone of hemolysis. It was thought that this streptococcus might be present quite generally on the mucous membrane in this patient. Cultures from the nose and pharynx and three out of six cultures from the stool proved that this was actually the case.

Sections of the excised muscle showed marked increase in the interstitial

tissue, poorly staining nuclei of adjacent muscle fibers, and slight round cell infiltration (Fig. 1). Sections of the fibrous nodule and fascia showed old and young connective tissue, absence of lymphoid tissue, small nests of round cells, plasma cells, and erythrocytes, chiefly around blood vessels. Gram-Weigert and methylene blue stains for bacteria revealed a moderate number of diplococci in or adjacent to the fibrous tissue between the muscle fibers and nests of cel-




Fig. 1.—Section of the muscle excised from the left side of the patient's neck. Note the marked infiltration by connective tissue, the irregular staining and the atrophy of muscle fibers. Hematoxylin and eosin. X 60.




Fig. 2.—Diplococci, singly and in short chains, and two leucocytes in tender fibrous node in the deep fascia from the left side of the neck. Methylene blue and eosin. X 1000.




Fig. 3.—Diplococci within the lumen and in the wall of the blood vessel and within the leucocyte just outside of the blood vessel in the center of the tender fibrous nodule of the deep fascia from the left side of the neck. Gram-Weigert. X 1000.

lular infiltration (Fig. 2) and a large number within and surrounding a small sized blood vessel in the center of the fibrous nodule (Fig. 3).

A vaccine was prepared by heating the streptococci suspended in salt solution from the primary cultures of the pulp of the tooth and muscle and treating it with equal parts of the patient's serum for two hours at 37°F and over night

in the ice chest. This was used in the treatment of the patient. The first dose consisted of 25,000,000, and was followed by marked muscle pains, especially at the left side of the neck, by nausea, extreme exhaustion and slight fever. The subsequent injections were at first diminished and then gradually increased, aiming to give a distinct but not severe reaction following the injection. Dr. Grimes, who referred the patient to me, reported three months after the extraction of the tooth, that the patient has shown gradual improvement; the attacks are milder and of shorter duration, the intervals between attacks are longer, and there has been marked improvement in the general nervous tone.

ANIMAL EXPERIMENTS.

Intravenous injections were made into rabbits, guinea-pigs and dogs, and intraperitoneal injections into mice. The streptococcus as isolated from the pulp of the tooth was injected into 4 rabbits, 2 guinea-pigs and 2 white mice, all of which recovered. The primary culture after one animal passage was injected into 2 rabbits, 1 dog and 2 mice. The rabbits and dog recovered; the mice died. Thirteen animals were injected. In 10 of these circumscribed hemorrhages and edema opposite the roots of teeth or at the foramina of exit of the superior or inferior maxillary nerves were easily visible; in 7 there were lesions of the muscles, 4 of which were limited largely to the muscles of the left side of the neck; in 6 there were lesions of the pulp of the teeth and the superior or inferior dental nerves; and in 3 there were insignificant lesions of the kidney, 2 in the gall bladder and 1 in the stomach. The spleen, adrenal, endocardium, vagus and sympathetic ganglia and subcutaneous nerves were normal. No lesions were found in the nerves supplying hemorrhagic muscles. The diphtheroid bacillus did not produce lesions in the one rabbit and one mouse injected.

The streptococcus from the muscle in the first and second cultures was injected into 2 rabbits, 1 guinea-pig and 1 mouse, producing lesions of the muscles, chiefly of the neck and shoulders, in all. In the rabbits there were also hemorrhage and infiltration of the dental nerves, and gross lesions in the dental pulps and periosteum opposite the roots of the teeth. In 1 there were a few hemorrhages of the stomach, in the other, hemorrhages of the kidney; in the mice there was a mild peritonitis.

To the portion of the above cultures of streptococci left over and having marked affinity for the muscles, was added 0.5 per cent formalin. This was allowed to stand over night at room temperature, a portion put aside for direct injection; the rest was centrifuged and the bacteria washed in Ringer's solution. The former was injected intravenously into a rabbit in amounts of 10 c.c. on four occasions over a period of eleven days. The rabbit showed a moderate number of small hemorrhages in the muscles of the hips and shoulders and a few in the stomach. The washed suspension was injected one month later into 3 rabbits, each receiving the growth from 150 c.c. of the broth cultures. All died in twenty-four hours. The muscles in all had a boiled appearance, and those about the shoulders, neck and spine had large and small hemorrhages associated with edema; 2 had, in addition, a hemorrhage of the pulp of the teeth; and 1 a few hemorrhages about the joints, in the tricuspid valve and in the left inferior dental nerve. The lungs were free from hemorrhages.

Two cubic centimeters or approximately 500,000,000 of the heat-killed streptococci failed to produce lesions, but this may have been due to the small size of the dose. The filtrate of the formalinized cultures failed to produce lesions in the muscles following 3 injections of 10 c.c. each into 1 rabbit. Three injections into 1 rabbit over a period of eleven days of the slightly hemolyzing streptococcus from the pharynx were followed by moderate numbers of lesions in the muscles in the right side of the neck, the right elbow and right shoulder and in the intercostal muscles. The streptococcus from the stools in the second cul-




Fig. 4.—Rabbit (R 656) showing marked lacrimation of the left eye and swelling of the left side of the face forty-eight hours after the intravenous injection of streptococcus from the pulp of the extracted first upper left molar.

ture did not produce lesions. The living streptococci from the tooth, muscle, pharynx and stool in the first or second cultures, and after one animal passage, and the heat or formalin-killed streptococci, were injected altogether into 24 animals (15 rabbits, 3 guinea pigs, 1 dog and 5 mice). Of these 17 (71 per cent) showed myositis, 13 (54 per cent) lesions of the periosteum opposite teeth or at nerve foramina 12 (50 per cent) lesions of the pulp of teeth and 10 (42 per cent) lesions of the dental nerves.² Similar results have been obtained since

²The incidence of the lesions in the other organs corresponded with those given in my paper on "Elective localization of streptococci," Jour. Am. Med. Assn., 1915, 45, 1687-1691.

in 2 cases of myositis and arthritis with streptococci isolated from the diseased pulp of extracted teeth.

Previous to these experiments lesions of the pulp of teeth and the dental nerves were not usually looked for, but recently careful search for similar lesions in these structures has been made in numerous animals injected with streptococci from other sources. In three instances only were there found lesions of the pulp of teeth and dental nerves. Two of these animals showed myositis in addition. The details in the following experiments will serve to illustrate specifically the results obtained:

R 656.—Belgian hare, 1510 grams weight. January 19, 1916, injected intravenously with the growth from 30 c.c. of ascites-dextrose-broth of a pure culture of streptococcus from the pulp of the extracted tooth.




Fig. 5.—Diplococcus in an area of hemorrhage in the left superior dental nerve of rabbit shown in Fig. 4. Gram-Weigert. X 1000.

January 20. Seemed quite ill; respirations were accelerated; appeared to have pain in walking and promptly crouched when quiet. The hair was roughened and there was marked lacrimation of the left eye, but no swelling of the face.

January 21. Seemed much better; was more active but tremulous, and appeared nervous. There was lacrimation of the left eye and an easily recognizable swelling of the left side of the face (Fig. 4). Slight pressure over this area appeared to cause pain and the swelling could be easily felt. Chloroformed. A rather large number of linear hemorrhages in the skeletal muscles, chiefly in the tendinous portions of the flat muscles of the shoulder and deeper muscles of the left side of the neck and of the front extremities were found. There were no lesions of the muscles of the hind extremities, the dorsal and lumbar regions of the spine, nor of the intercostal muscles and diaphragm. On removing the skin on the left side of the face, marked edema, infiltration and hemorrhage of the subcutaneous tissue, the fascia, the muscles and the periosteum, were found. The hemorrhages in the periosteum opposite the molars appeared to be the center of the edematous area. The hemorrhagic infiltration extended to the under surface of the orbit. The pulp of the left third and fourth upper molars was found to be edematous and hemorrhagic. The hemorrhages were small and punctate and were not found in the pulp of the 2 corresponding teeth on the

opposite side. On dissecting away the soft tissues of the lower jaw, it was found that hemorrhages of the periosteum had occurred opposite the apices of the right lower incisor, and the first and two last molars, the second left lower molar, and surrounding the right mental foramen similarly to the lesions shown in Fig. 6. The pulp of the teeth surrounded by hemorrhages in the periosteum was found to be edematous and hemorrhagic, whereas 3 adjacent pulps appeared




Fig. 6.—Left lower jaw of rabbit (R 659) injected three days previously with 5 c.c. of twenty-four hour tissue-ascites-fluid culture of the streptococcus from the tooth. Note the edema and hemorrhage in the periosteum opposite the root of the canine and surrounding the inferior dental nerve at the foramen of exit. X 2.



Fig. 7.—The pulp of the inferior canine and inferior dental nerve from dog (D 412) two days after an intravenous injection of the streptococcus isolated from the hemorrhagic pulp of the second upper left molar. Note the numerous large and small hemorrhages in the pulp of the left lower pulp and nerve. X 2.

normal. The left superior dental nerve was extremely hyperemic and a number of small hemorrhages were found in the sheath. The only lesions of the viscera were 3 small hemorrhages in the mucous membrane of the pyloric end of the stomach and 3 small subserous hemorrhages at the fundus of the gallbladder.

January 23. Blood-agar plate cultures of the blood produced 1 colony of streptococcus; of the emulsions of the pulp of teeth 5 and 18 colonies respectively. Cultures in ascites-dextrose broth from the emulsions of the pulp of 2 teeth, of

the hemorrhagic muscle, of the edematous tissue over the jaw and of the hemorrhagic periosteum, showed a short chained streptococcus.

Sections of the edematous tissue over the left jaw showed extreme hemorrhage, edema and beginning leucocytic infiltration. In the pulp of 2 teeth from which sections were made, were large and small hemorrhages chiefly beneath the layer of odontoblasts, and in the left superior dental nerve were small areas of hemorrhage in the sheath. Sections of hemorrhagic areas in the scapular muscles showed marked hemorrhage between the muscle fibers, separation and fragmentation of the latter, and slight leucocytic infiltration. Gram-Weigert stains showed diplococci in or adjacent to the hemorrhagic areas in the edematous periosteum, in the muscle from the scapula, in the pulp of the teeth, and in the left superior dental nerve (Fig. 5).

D 412.—A white and black dog, weight 9.5 kilos. January 22, injected intravenously with the growth from 90 c.c. of ascites-dextrose broth of the streptococcus from the hemorrhagic pulp of a tooth of the above rabbit.




Fig. 8.—Section of the pulp of the left lower canine shown in Fig. 7. Note the marked hyperemia and the large and small hemorrhages. Hematoxylin and eosin. X 60.

January 24. Seemed well; was active; no swelling of the face; no tenderness over joints nor muscles. Chloroformed. The left inferior dental nerve was found to be edematous and hyperemic and contained a number of large and small punctate hemorrhages (Fig. 7). The pulp of the corresponding canine tooth likewise appeared edematous and contained numerous punctate hemorrhages. The corresponding nerve and pulp on the opposite side appeared normal (Fig. 7). There were similar but less marked lesions of the left superior dental nerve and pulp of the canine tooth. The first upper and lower molars were examined and only the first left lower molar showed unmistakable hemorrhage. The muscles were free from visible lesions except for a few hemorrhages in the flat muscles under the scapulæ. In a painstaking examination of all the organs no other apparent lesions were revealed.

January 25. Blood-agar slants made from the blood were sterile.

Sections for microscopic study were made from the left inferior dental nerve, from the pulp of the teeth mentioned, and from the superior maxillary

nerve. Hemorrhages and beginning leucocytic infiltration were found in the pulp of the left inferior canine (Fig. 8) and a small number of hemorrhages in the pulp of the first left lower molar, but practically none in the others. These hemorrhages were most numerous immediately beneath the layer of odontoblasts and at the distal portion of the pulp (Fig. 8). Sections of the left inferior dental nerve showed marked edema, a moderate leucocytic infiltration and a number of large and small hemorrhages chiefly in the sheath (Fig. 10). There



Fig. 9.—Diplococci, singly and in short chains, within and outside leucocytes, adjacent to the area of hemorrhage shown at *a* in Fig. 8.



Fig. 11.—Diplococci and double chain of streptococci in the hemorrhagic and infiltrated area (*a*) in the sheath, shown in Fig. 10.




Fig. 10.—Section through the hemorrhagic area in the left inferior dental nerve, shown in Fig. 7. Note the marked edema and large and small hemorrhages and leucocytic infiltration of the sheath. Hematoxylin and eosin. X 60.

were no lesions in the left superior maxillary nerve. Gram-Weigert and methylene blue stains for bacteria showed scattered diplococci and at times chains of diplococci in and adjoining the hemorrhagic areas (Figs. 9 and 11). In one instance, a mass of diplococci surrounded by erythrocytes was found just outside the wall, and 2 diplococci in the wall of a small blood vessel. In a number of instances the diplococci were found within leucocytes and in what appeared to be endothelial cells. No bacteria could be found in the normal portions of the pulps and nerves showing lesions, nor in those free from lesions.

P 155.—White guinea-pig, weight 320 grams. February 19, injected intravenously with the growth from 30 c.c. of ascites-dextrose broth of the streptococcus isolated from the muscle of the patient.

February 21. Seemed quite well, but appeared to be muscle-sore. Chloroformed. Numerous small hemorrhages associated with edema in the triceps muscles and a moderate number in the muscles of the left shoulder and a few hemorrhages in the deeper layer of the muscles of the left side of the neck




Fig. 12.—Section of the dental pulp of the left upper molar in guinea pig (P 141) injected two days previously with the streptococcus from the pulp of the patient's tooth. Note the marked leucocytic infiltration and hemorrhage. Hematoxylin and eosin. X 240.




Fig. 13.—Diplococcus in infiltrated area (a) shown in Fig. 12. Gram-Weigert. X 1000.

were found. There were no lesions of the dental nerves but the pulp of the two left upper molars was extremely hyperemic and contained distinct hemorrhages. There were no other findable lesions.

February 24.—Cultures from the blood in ascites-dextrose broth were sterile.

Sections through the areas of hemorrhage in the scapular muscle showed marked extravasation of blood corpuscles, leucocytic infiltration, separation and necrosis of muscle fibers (see Fig. 14 illustrating similar lesion in triceps muscle).

Sections of the pulp of the left first upper molar showed a number of small hemorrhages while in those of the superior maxillary and right superior dental nerve there were no lesions. Gram-Weigert and methylene blue and eosin stains showed a few scattered diplococci in the hemorrhagic and infiltrated area in the muscle and adjacent to the hemorrhages in the tooth-pulp (Fig. 15).



Fig. 14.—Section of the left triceps of guinea pig (P 155) injected two days previously with the streptococcus from the muscle of the patient. Note the marked hemorrhagic and leucocytic infiltration and separation of muscle fibers. Methylene blue and eosin. X 140.



Fig. 15.—Diplococcus in infiltrated area shown in Fig. 14. Gram-Weigert. X 1000.

THE STREPTOCOCCUS.

The streptococcus isolated from the dead pulp of the tooth, the muscle, the pharynx, and the stool, produced small, round, slightly elevated grayish-brown, non-adherent, rather dry, colonies on aerobic blood (human) agar plates. In broth it produced a diffuse turbidity with a small amount of flocculent sediment at the end of forty-eight hours. It acidified and coagulated milk. In liquid media it grew in short chains and diplococci. In many instances, the single members of the diplococci were quite round and resembled staphylococci. The strains from the tooth and muscle were freely susceptible to phagocytosis

and the virulence was of a low order. The streptococci from blood agar slants in salt solution used as a vaccine resisted 60°C. for thirty minutes on consecutive days, while after the third heating the subcultures remained sterile. The blood of the animals soon became sterile, and nearly all recovered promptly from the effects of injection. The organism resembled very closely the streptococci that



Fig. 16.—Section of the left trapezius in rabbit (R 660) forty-eight hours after intravenous injection of the streptococcus from pulp of the left upper molar in rabbit shown in Fig. 4. Note the marked edema, leucocytic infiltration, hemorrhages and the disintegration of the muscle fibers. Hematoxylin and eosin. X 120.

I have isolated from other cases of myositis and strains from other sources having affinity for muscles.

In Table 1 is given the fermentative power of the strains isolated from the pulp of the tooth, the muscle, the pharynx, and the stool. It is seen here, as has been found to be the case with other streptococci, that there is little parallelism between pathogenicity and the degree or range of fermentative powers.

TABLE 1
FERMENTATIVE POWERS OF THE STREPTOCOCCUS*

Streptococcus	Dextrose	Lactose	Saccharose	Maltose	Raffinose	Mannite	Salicin	Inulin
Tooth pulp.....	2.4	1.5	0.9	1.9	0	1.3	2.0	0
Pharynx.....	1.9	1.8	0	1.6	1.8	1.0	0	0
Muscle.....	1.5	1.2	1.5	1.3	0	0	0	0
Stool.....	2.9	2.3	1.1	3.2	0	0	2.3	0

*The standard sugar-free broth containing the usual amounts of the various carbohydrates was used in the fermentative tests. The cultures were incubated 72 hours. The figures indicate the number of cubic centimeters of normal tenth sodium hydrate required to neutralize 5 c.c. of the broth culture, phenolphthalein being used as an indicator.

SUMMARY.

A streptococcus having peculiar properties was isolated from the dead pulp of the left upper first molar in the region where the attacks of pain usually began. The streptococcus was also demonstrated in the sections and isolated from the

infiltrated deep fascia and muscles of the left side of the neck. A similar streptococcus was isolated from the pharynx and stool. This streptococcus was proved to have elective affinity for the pulp of teeth, dental nerves and muscles in animals. It was repeatedly isolated from and demonstrated in the experimental lesions in animals whose blood was sterile; the lesions were again produced on reinjection and the streptococcus again isolated. Many animals appeared to be in pain and 1 rabbit (Fig. 4) had marked swelling and tenderness over the left upper jaw. This affinity was proved absent in the diphtheroid bacillus and *Bacillus fusiformis*, which also were isolated from the pulp of the tooth, and in the streptococcus broth culture filtrate. Streptococci from other sources rarely cause lesions in the pulp of teeth and dental nerves. The phagocytic power of the patient's blood following the attack over the strain from the tooth was twice that of comparable normal blood.

These results would appear to warrant drawing the conclusion that the attacks of pain in the face in this patient were due to a streptococcus infection of the sheaths of the dental nerves, and that the pain, swelling, tenderness and spasm of the muscles of the neck were due to myositis and fibrositis—the result of infection by this streptococcus. The demonstration of living streptococci in the pulp of the tooth and in the fascia and muscle during quiescent intervals is significant and may explain the recurrence of the attacks. The cavity in the tooth containing the dead pulp, which was originally infected from the mouth, judging by the character of the filling and of the bacterial flora, was quite unable to heal for mechanical reasons. The contents of the cavity appeared to afford a culture medium for the growth of the streptococcus. From stimulation of the defensive mechanism in the patient during the attacks, active growth appeared to be held in check and the symptoms disappeared in consequence, only to reappear later from recurrence of active growth and localization of the streptococci when the immunity was low.

The improvement in the patient since extraction of the tooth appears to be due to the removal of this focus and to prolonged artificial stimulation of the defensive mechanism by means of the autogenous vaccine, which, it is hoped, will lead to the destruction of all the streptococci in the muscle and dental nerves, and result in the ultimate recovery of the patient. However, the isolation of this streptococcus from so many places, indicates that complete recovery will probably be difficult.

PLEA FOR A MORE GENERAL USE OF THE X-RAY

BY DR. HAROLD B. THOMPSON, SEATTLE, WASH.

THE value of x-ray in the diagnosis of dental pathology has been appreciated by a few men in both the dental and medical profession for a number of years. More recently the increase of x-ray equipment, together with considerable literature on the subject, both in professional and lay periodicals, has wonderfully broadened the field and increased the application of this modern diagnostic aid.

There are still a large number of men in both the dental and medical fraternities who do not avail themselves of this help in diagnosis, at least to a sufficient extent for their own welfare. Whether this neglect is due to ignorance, habit, or an exaggerated opinion of one's own diagnostic ability, it is certainly inexcusable. There still remain a few physicians who will volunteer to treat fractures without the aid of the x-ray, even when radiographs are available; but through slow education, the use of the x-ray has become so universal, that the failure to procure an x-ray examination in fractures, when possible, now constitutes a *prima facie* cause for an action of malpractice. It is only a question of time, if the time has not already arrived, when the failure of the dentist or physician to avail himself of the aid of the x-ray in dental conditions, will render him liable for malpractice to the same extent. The use of the x-ray then, constitutes a form of insurance against malpractice suits, and the premium is paid by the patient.

The work of Dr. Rosenow along the line of bacterial origin of arthritis, ulceration, appendicitis, gall bladder diseases, etc., is familiar to all in the profession. So much has appeared in the lay literature recently along this line, that the public is beginning to demand an x-ray examination of the teeth, sometimes even before the physicians or dentists. Although all my work is referred work, I have recently had several cases appear for dental radiographs without being referred, when the symptoms were confined to general joint conditions or to the abdominal organs. It is certainly more to the credit of the attending man to anticipate a patient's wishes in this regard than to have the course of action suggested by the patient.

It is often a desire for economy on the part of the attendant when he does not ask or insist on a radiograph. Numerous experiences have shown me the fallacy of this line of procedure. One dissatisfied patient can do more harm than several pleased ones can do good. A patient who has been treated for several weeks or even months by one man and then changes to another, who refers the case for a radiograph and with its aid completes a cure in a short time, has good reason to become a "knocker." A patient's health is his most valued possession, and he will appreciate your interest in his case as shown by your desire for all the information possible, even though the radiograph gives no material aid.

I have intentionally omitted in the body of this article, an enumeration



Fig. 1-A.



Fig. 1-B.

Fig. 1.—Typical apical abscesses.



Fig. 2.—Absorption of tip of root and decalcification of surrounding alveolar process.



Fig. 3.—Small area of rarefaction (apical abscess) at tip of lateral incisor.



Fig. 4.—Small areas of rarefaction at tips of right middle and lateral incisors.



Fig. 5.—Small fragment of alveolar process left after extraction of tooth and causing persistent sinus.

of the specific class of cases in which the x-ray has proved of inestimable value, as this has been ably covered time and again in recent literature. My plea is more for a general line of attack, than a plea for the use of the x-ray in any specific class of cases, and you will find this general attitude of availing yourself of all possible aids in diagnosis to rebound to your benefit.



Fig. 6.—Decalcification of alveolar process surrounding root of crowned pivotal tooth from pyorrhea.



Fig. 7.—Decalcification at surface of alveolar process, extending downward at side of canine and appearing also at tips of middle and lateral incisors, pyorrhea.



Fig. 8.—Unerupted canine.



Fig. 9.—Impacted third molar causing "neuralgia."



Fig. 10.—Unerupted canine causing "neuralgia" by pressure against the root of the lateral incisor.



Fig. 11.—Incomplete filling of root canal.

A recent case which came under my observation, shows the value of another laboratory test, but the application is the same. A specialist sent a case to the hospital one day and engaged the operating room for the following morning, intending to do an extraction of all the teeth for an aggravated case of pyorrhea. The young interne at the hospital noticed the peculiar pallor of the patient and



Fig. 12.—Showing acute curve in root, giving excuse for drilling through side of tooth.



Fig. 13.—Hole drilled through side of tooth with filling protruding, causing necrosis of alveolar process.



Fig. 14.—The dentist has drilled through the side of the root and then inserted a root canal filling which has caused necrosis of the bone. In attempting to remove this filling a small pair of forceps has been broken off outside the tooth.



Fig. 15.—Protrusion of root canal filling, causing necrosis.



Fig. 16-A.



Fig. 16-B.

Fig. 16.—Stereoscopic radiographs showing the direction of growth of unerupted teeth.

took a blood smear, which showed an advanced typical case of leukemia. Needless to say the operation was postponed, and well it was, for the patient died within a week, and his death would probably have been attributed to the operation, had it been done.

The accompanying illustrations will serve to call to mind some of the more obvious specific cases in which great benefit may be expected from the use of the x-ray.

My plea for a more general use of all possible aids in diagnosis, particularly that of the x-ray, as it has a much greater field in dental work than any other laboratory method, may be summarized as follows:

1. A more general use of the x-ray will allow a more positive diagnosis with consequent more particularized treatment, and better and quicker results.
2. The more general use of the x-ray is a form of insurance against mal-practice suits with the premium paid by the patient.
3. The risk of having your patient become a "knocker" is transferred to the man who fails to use the x-ray.
4. The appreciation of your thoroughness when the patient is convinced that you are using every means at your disposal to alleviate his condition, will prevent dissatisfaction in many cases and engender a confidence which even unfortunate results will not be able to overcome.

News and Notes

Dr. E. Santley Butler, practice devoted exclusively to orthodontia, has announced removal of offices from Room 704 to Suite 605, 576 Fifth Avenue, New York City.

* * * * *

Drs. Sturdevant and Dinham, practice limited to orthodontia, announce their location at 919-920 Selling Building, Portland, Oregon.

* * * * *

Dr. Blaine Truesdell announced the removal of his offices to Suite 1112 First National Bank Building, Omaha, Nebraska. Practice limited to orthodontia.

The International Journal of Orthodontia

PUBLISHED THE FIFTEENTH OF EVERY MONTH BY

THE C. V. MOSBY CO., 801-807 Metropolitan Bldg., St. Louis, Mo.

Foreign Depots—*Great Britain*—Henry Kimp-ton, 263 High Holborn, London, W. C.; *Australasia*—Stirling & Co., 317 Collins Street, Modern Chambers, Melbourne; *India*—"Practical Medicine," Egerton Street, Delhi; *Porto Rico*—Pedro C. Timothee, Rafael Cordero 68, San Juan, P. R.

Subscription Rates—Single Copies, 30 cents. To anywhere in United States, Cuba, Porto Rico, Canal Zone, Mexico, Hawaii and Philippine Islands, \$3.00 per year in advance. Under foreign postage, \$3.40. English price: 15/ per annum, 1/6 per number. Volume begins with January and ends with December of each year.

Remittances—Remittances for subscriptions should be made by check, draft, postoffice or express money order, or registered letter, payable to the publishers, The C. V. Mosby Company.

Contributions—The editor will be pleased to consider the publication of original communications of merit on orthodontic and allied subjects, which must be contributed solely to this journal.

Opinions—Neither the editor nor the publisher hold themselves responsible for the opinions of contributors, nor are they responsible for other than editorial statements.

Reprints—Since it is not desirable to hold type standing longer than absolutely necessary, all requests for reprints should be made at time of submitting manuscript for publication. Rate card will be sent with galley proof.

Communications—Contributed articles, illustrations, letters, books for review, and all other matter pertaining to the editorial department should be addressed to the Editor, Doctor Martin Dewey, 1016 East Armour Boulevard, Kansas City, Mo. All communications in regard to advertising, subscriptions, change of address, etc., should be addressed to the publishers, The C. V. Mosby Company, 801-807 Metropolitan Building, St. Louis, Mo.

Illustrations—Such halftones and zinc etchings as in the judgment of the editor are necessary to illustrate articles will be furnished when photographs or drawings are supplied by the authors of said articles.

Advertisements—Objectionable advertisements will not be accepted for publication in this journal. Forms close first of month preceding date of issue. Advertising rates and sizes on application.

Change of Address—The publishers should be advised of change of subscriber's address about fifteen days before date of issue, with both new and old addresses given.

Nonreceipt of Copies—Complaints for nonreceipt of copies or requests for extra numbers must be received on or before the fifteenth of the month of publication; otherwise the supply is apt to be exhausted.

Entered at the Post Office at St. Louis, Mo., as Second-Class Matter

EDITORIALS

The Universal Regulating Appliance

REGULATING appliances are mechanical devices for exerting force on the malposed tooth for the purpose of creating cell activities thereby causing the tooth to assume its proper position in the line of occlusion. A universal regulating appliance is one that is capable of being used to the best advantage upon all cases. The ideal regulating appliance must possess certain requirements, the first of which is efficiency. It must be capable of doing the things which it is intended to do. It must exert force upon the malposed tooth with the result that cell activity will occur and the tooth be moved into the line of occlusion. After efficiency, there are other things to be taken into consideration with reference to making an ideal appliance. One of the requirements of any appliance is that of stability and anchorage, by which we mean the appliance must be so attached as to be capable of exerting the desired force. It must also be so attached that it will exert force in the proper direction so as to produce the proper cell activity and tooth movement.

In going through orthodontic literature we find that there are several appliances which seem to approach the ideal. There are several styles of appliances upon the market which, in the hands of certain men, will easily accomplish the desired result. These appliances may be divided into two groups, known as "fixed" and "removable," each one of which has certain advantages and disadvantages. When we consider the radical mechanical difference between those two appliances it seems almost an impossibility to talk about any one appliance which will fulfill the requirements possessed by both the "fixed" and "removable." It has been stated by Dr. Eby that the regulating appliance of the future will be one that will embody the advantages of both the fixed and removable, combining them into an appliance which might be termed the "fixed removable" type. In studying the results of orthodontic treatments of times past, we might say that so far as mechanism and the construction of an appliance in regard to tooth movement is concerned, we have appliances which are ideal. We have several styles of appliances which seem to be able to correct any case of malocclusion encountered. In other words, if the movement of the malposed tooth was the only thing to be considered we would be able to select several styles and kinds of appliances, any of which would produce that movement. However, the movement of the malposed tooth is not the only thing to be desired as has been revealed to the sorrow of some men. In considering an ideal regulating appliance, we have to consider the material from which the appliance is constructed. When we take up the question of material we are immediately confronted with one of the biggest problems we have to contend with from the standpoint of ideal conditions. Up to the present time we have no metals or materials which may be considered ideal. A metal which possesses one requirement to a very high degree of efficiency will be decidedly lacking in another. For instance, several years ago the majority of regulating appliances on the market were made out of German silver. German silver possesses a great many ideal characteristics, the principal one of which is the ease with which it may be worked and also its antiseptic properties. Looking over the large number of cases which have been treated with German silver appliances in the past, we find some very nice results have been obtained; we also find some of evil effects. We find that in a great many instances some of the anchor teeth were discolored, were turned black from a black stain or oxide very difficult to remove. In checking up these results, however, we are forced to admit the actual amount of decay which occurred in the mouths of those treated with German silver appliances was comparatively slight. However, as a result of the tarnishing of the German silver and as a result of its appearance, it was not satisfactory from an esthetic standpoint. Therefore, a large number of men began using gold and platinum and iridio-platinum, because the appliances were more pleasing in appearance. In other words, they sacrificed the antiseptic properties in the metal for the esthetic, with the result that an examination of the cases treated by gold and platinum and iridio-platinum appliances show a very great loss of tooth structure. In other words, the amount of decay which has occurred with the use of the iridio-platinum and gold and platinum appliances is far in excess of that which occurred in the use of the German silver appliance. To eliminate this decay which resulted from gold and platinum appliances, some men began advocating the banding of every tooth that the

appliance would touch, advocating a layer of cement to protect the teeth. The use of so many bands on the teeth to prevent them from decaying defeated the purpose of the better appearance of the noble metals as compared to German silver, for nothing is more unsightly than a large number of bands on anterior teeth.

In using bands there is always the ultimate danger of the bands resting against the tissue, and we now have oral prophylactic specialists who condemn orthodontic treatment because of the injuries to the gingival tissues produced by bands. The facts remain that if one band occasionally injures the gingival tissue, that injury is more apt to occur where a large number of bands are used than if very few bands were used. Even in the construction of the removable appliance which does not employ any bands, we have to take particular care to avoid injury of the gingival tissue with the gingival extension which is made to go under the lingual convexity to hold the appliance; but there is very little liability to decay providing the appliance is carefully cleaned and taken care of as it should be. However, there is one disadvantage in the removable appliance,—the placing of the appliances on the lingual side may interfere with the speech of the patient. Therefore in this respect the removable appliance is not ideal.

We have mentioned that appliances were changed from German silver, to gold and platinum, and iridio-platinum for the esthetic effect, which brings out the point that the esthetic effect or appearance of the appliance is an important thing to be considered. A great many patients do not care how efficient the appliance is from a mechanical standpoint, if it looks well, and they prefer a less efficient appliance to one that is conspicuous. Therefore we have to consider the question of appearance in regard to regulating appliances, and the ideal appliance must be so constructed as to be inconspicuous if not entirely out of sight. In regard to inconspicuousness, the majority of removable appliances have an advantage over a large number of fixed appliances.

Another requirement of the ideal appliance, about which many statements have been made is that it must produce physiological cell activities. Just what the activities of cells are as a result of the regulating appliance is not exactly understood. We do not know, however, that the results of some of these appliances have not been satisfactory from the standpoint of cell activity. We have known of several cases where the roots of the teeth have absorbed for some reason or other during tooth movement. We do not know exactly why this was, except as a general statement, that the proper cell activity was not produced. The question remains, "What is the proper cell activity to be expected from regulating appliances?" The answer to this question can only be given in a general statement: namely, that the cell activities produced by the regulating appliance should be in keeping with the same cell activity that occurs as a result of normal development. In other words, malposed teeth moved by regulating appliances should follow as nearly as possible the movement produced by a tooth during the process of eruption. To approach this style of tooth movement, the appliance must be so constructed that the teeth will be free to respond to the forces of mastication and not lashed or bound rigidly by any fixed or firm appliance. Also during the eruption of the majority of teeth they do not travel in a straight line, they do not travel bodily in one direction, but tip more or less during their movement. If you will examine the position of the upper teeth

before eruption, as they lay in their crypts, it will be seen that they must travel buccally a considerable distance and a large amount of that traveling is the result of tipping. The lower molars and premolars as a rule travel lingually to a certain extent which also is the result of tipping. This lingual movement of the lower molars is especially true in the deciduous teeth, although the permanent teeth do not travel lingually as much as the deciduous teeth. Any appliance then that exerts pressure to produce physiological cell activity will be one that will not interfere with the individual development or growth of any one tooth, neither will there be undue pressure brought to bear on the teeth in any one particular direction.

We have outlined a number of the requirements of the ideal appliance, and in a general way, stated some of the difficulties which are encountered. With the large variety of malocclusions which we are called upon to treat at the present time it seems that the best plan is not to look for a universal regulating appliance, but choose the appliance which more nearly approaches the ideal in that particular case. An appliance which may be desirable in one individual or case of malocclusion would not be so ideal in another case. Therefore, we believe at the present time, that the greatest service will be rendered the patient if the orthodontist will become able to analyze regulating appliances from a mechanical standpoint, and consider what is required in the treatment of the particular case in hand and select the appliance from the standard types which have been proved worthy and useful. We do not believe that it is necessary to make and design a separate appliance, based on new mechanical principles for every case of malocclusion which we treat. Any regulating appliance regardless of whether it is a fixed or removable variety exerts pressure upon the teeth in only two ways, either pushing and pulling, or by a combination of the two, which is twisting. After an appliance is secured which does that, there only remains the selecting of an appliance with some means of properly attaching it to the malposed tooth to obtain the desired force to produce the mechanical stimulation which will cause the cells to respond in a physiological manner. We also believe it is a mistake to say any one style of appliance, because it has been advocated by a group of men, is the only kind to use. We must realize that in certain instances other appliances might better serve the need of the case, be more comfortable to the patient, and be less conspicuous and more hygienic. There are even instances when a simple appliance will do the work as satisfactorily as a more complicated one.

Therefore, the question of the regulating appliance of the future remains for the individual operator to be able to analyze his case so as to be able to select the best appliance for the particular case; and possibly, far in the future, we may evolve an ideal regulating appliance.

Is Our Dental Literature Free from Commercial Influence?

THERE is one question which arises in the comparison of the literature of the dental profession with that of the medical profession and other scientific bodies. That question is whether our dental literature is controlled or influenced by commercial houses, or whether commercial influence is exerted upon the

publication of our articles; for the majority of our dental journals, those which have the largest circulation and which may be said to be representatives of dental literature, are journals controlled by trade houses. It is a lamentable fact that the dental profession, at the present time, has but few journals that are not influenced by trade houses or under the control of houses interested in selling dental goods.

We are aware of the fact that the medical profession passed through a similar stage several years ago, before the *Journal of the American Medical Association* reached its present standing. This resulted in the improvement of medical journals controlled by publishing houses, and the elimination of trade journals. We now have high class scientific journals which have no commercial appliances for sale and which also are free from, or not interested in, the selling of medical supplies.

We have often wondered whether the dental journal that is controlled and published by a house whose sole object is the selling of dental supplies is published in the interest of the profession, or whether it is not published for the advertising of their own products. We are aware of the fact that editors of the dental journals have very high ideals, and, so far as the ideals of the editors, or the editors themselves, are concerned, we have no criticism. However, it has been brought to our attention that several of the editors of dental journals are heavy stock-holders in the supply houses that publish journals. We, therefore, wonder if those particular editors could serve the interests of the profession as well as if they were not interested in the selling of dental supplies. We wonder if they would publish in their journals papers written by scientific men criticizing very severely some of the articles in the sale of which the supply house is interested, and which are advertised very extensively in the advertising department of the journal. It might be that our dental editors would wish to do so, but we very seriously doubt that the supply house or the publishers of the journal would allow such a thing to occur. It has been shown that there are several very extensively advertised cements on the market which, by scientific analysis, do not carry out the claims made by their manufacturers. We, therefore, wonder if the publishers would allow a criticism of such cements from a purely scientific standpoint to appear in their journal at the same time their advertising pages would be highly lauding the advantages and wonderful properties which they possess. We believe that the dental editor would be more able to serve the interest of the profession from a scientific standpoint if his sole business was the editing of the journal, and if he was not interested in the sale of dental articles.

In some journals, articles are published as scientific which are really only advertisements for some material or appliance. Papers are also often written for the purpose of selling articles which are advertised in the pages of the journal. We remember recently a very prominent dental trade journal which contained a featured paper on the first pages describing a new appliance. This particular appliance may have a great many virtues, but it is often plain to see that some papers are written to sell appliances. A great many claims are made for that purpose. In the same issue of that trade journal there appeared a featured advertisement calling attention to the wonderful qualities of an appli-

ance and referring back to a paper published as a scientific paper. Since that time the supply house has sent out advertisements broadcast in which they include reprints of the paper, which are mailed along with the catalogue of the appliance. The advertisements of the appliance are constantly referring to the paper which was published in the journal. Would the same prominence be given to a paper which presented scientific criticism of the aforesaid appliance? Would they publish a criticism of an article which they widely advertised? We believe that commercialization of scientific papers from a professional standpoint is very poor policy. If papers were written showing the weakness of widely advertised articles, would the publishers be as willing, for the scientific advancement of orthodontia, to give a paper criticizing some of their appliances the same publicity they would give the paper calling attention to the virtues of the appliance. We believe that the virtues of any article should be published broadcast, but we also believe that it should be published in a scientific manner and not in a commercialized manner.

If dentistry is a science, it must have journals that are willing to publish scientific papers even if the papers criticize materials and appliances that are products of the publisher. It has been called to our attention that there are some trade journals that will not illustrate any style of an appliance unless manufactured by their own house. This again is a commercialization of scientific literature which handicaps the profession, and which the dental profession should remember, contributes to the weakness of their literature as compared with that of the medical profession.

The Small Gauge Regulating Appliance

SEVERAL years ago the majority of appliances or alignment wires placed upon the market were made of 16 gauge material. The 16 gauge arch was the universal regulating appliance which at that time was made principally of German silver or some other alloy. When it became the common plan of treatment to correct malocclusion of the deciduous teeth, the 16 gauge arch was found to be too large. There then was a tendency towards the use of 18 gauge arches and at that time the 18 gauge was the smallest arch placed upon the market. Since that time arches have been manufactured in all gauges varying from 16 to 20 and some even as small as 36 thousandths of an inch. As a result of the use of the small gauge arch, some men have been inclined to discard the large gauge in favor of the small, claiming the small gauge is less annoying to the patient and consequently should be used in all cases. We are willing to agree that the small gauge arch up to 19 or 20, or even 25 or 36 thousandths of an inch, may have advantages in some cases; but, it by no means replaces the large arch, in some cases, if the most efficient regulating appliance is desired. The small gauge arch has its limitations as well as its usefulness and it should by no means be considered the ideal appliance for all conditions.

The small gauge arch can be used satisfactorily in those cases of malocclusion which require a reciprocal anchorage, in those cases of neutroclusion where all the teeth have to be expanded an equal amount; it is ideal because of

its delicate spring which works for a considerable length of time without readjusting. We find in other types of malocclusion that it possesses several disadvantages. One disadvantage of the small gauge arch is that it must be fitted very close to the teeth, for owing to its small diameter, it is much more liable to injure the cheek than an arch of a larger size. If the patient lies on the side of his face during the night the small gauge arch is more liable to cut into the cheek than a 16 gauge arch. Owing to the small gauge arch possessing a large amount of spring it is contraindicated in those cases where the point of attachment is far removed from the anchorage. In other words, if the small gauge is used on cases where it is necessary to move the four anterior teeth forward without any expansion in the molars or premolars, it possesses disadvantages because of the tendency to spring buccally between the attachment on the incisors and the molar anchorage. The same trouble would recur if the small gauge arch of 25 or 36 thousandths of an inch was being used to open up the space for a premolar by carrying the anterior teeth forward by the action of the nut. If we were going to have stationary anchorage on one side and primary anchorage on the other, or if our case was one that required expansion in the molar region on one side and the holding of the molar stationary on the other, it would be difficult to secure stationary anchorage with the small gauge arch owing to the elasticity which would permit the arch to spring. Likewise, in the use of the intermaxillary anchorage where we desire the backward movement of the teeth, as in the treatment of a mesiocclusion case, there is a great tendency for a small gauge arch to spring between the molar anchorage and the attachment of the intermaxillary rubber. These are only a few of the conditions where a small gauge arch is contraindicated, but for the benefit of a large number of men, especially the younger men, who may become over enthusiastic over the use of the small gauge arch, we are mentioning these disadvantages. Dr. Lourie has stated that the shorter the regulating appliance between the points of attachment, the smaller the gauge can be. In other words, there is a limit to the smallness of the gauge of the alignment wire where it is lingual or labial, and this limit depends upon the nature of the teeth movement to be accomplished, the direction from which the force is to come from the appliance, whether it is to be entirely a spring force or a pushing force derived from the action of the nuts, whether the attachments to the appliance are close together, and whether reciprocal anchorage or stationary anchorage is to be used.

The advantages and disadvantages of the small gauge arch should be studied from a scientific standpoint, and it will be found useful in one case and a hindrance in another.

Dr. William Goode

DR. WILLIAM A. GOODE, Orthodontist, San Diego, California, was drowned in the Cuyamaca Lake, Sunday, November 19th, when his light duck boat was capsized by a strong wind. Dr. Goode with a number of San Diego physicians and dentists had gone to the lake Saturday night for an outing over Sunday. Dr. Kerch and Dr. Goode were in a light duck boat which

was capsized by a large wave about two hundred yards from shore. After clinging to the boat a short time, Dr. Goode, who was very athletic, started to swim to shore. He had gone about half way when he disappeared, probably as a result of the chilly water and also being weighed down with heavy hunting clothes and ammunition. Dr. Kerch was rescued by hunters in another boat, and they attempted to reach Dr. Goode when he sank.

Dr. Goode was thirty-four years of age, having lived in San Diego twenty-two years. He was a graduate of the Sherman Heights School, the San Diego High School, and Pomona. While a student at Pomona, he was a member of the Pomona football team. He graduated from the Northwestern University Dental School in 1910, and took a postgraduate course in the Dewey School of Orthodontia.

Dr. Goode leaves a widow, who, before her marriage, was Miss Ethel Speaker, physical director of the San Diego High School.

Dr. Goode had a very brilliant future, he was very well esteemed by both the medical and dental professions. While he had only shortly taken up the practice of orthodontia, from the success he obtained in other lines, we are sure that orthodontia loses a good man by the untimely death of Dr. Goode.

A New Fourth Edition of Prinz "Dental Materia Medica and Therapeutics"

THE dental profession is now happily informed of Dr. Hermann Prinz's new (4th) Edition* of "Dental Materia Medica and Therapeutics," and is hereby potentially offered a new cause of congratulating itself upon the numerous enrichments presented in the new volume. While our dental literature may be apparently well supplied with text-books treating the subject either as a separate branch or in connection with dental operative procedure which might well answer the demands of the student and practitioner, yet a new volume is always welcome if it exhibits some special features or "strong points" not already elaborated in similar works. The many new and special features in Prinz's new book meet our eyes by first impression upon casually glancing through its pages, and upon closer analysis, we find the new volume is as fairly concise as it is fully comprehensive. The author's indefatigable efforts to establish dental therapeutics on modern pharmacological research within the fullest limits of present possibilities must finally win our admiration for his strong advocacy for rationalism as against empiricism. We may define materia medica and place as a synonym with it the term "pharmacology" if we hold to this distinction that the latter name, in addition, implies the *specific action* of the drugs upon the various tissues, organs, and functions of the body. In this sense, Dr. Prinz has given us more than a materia medica, he has presented us with a dental pharmacology of inestimable value.

*Dental Materia Medica and Therapeutics. By Hermann Prinz, D.D.S., A.M., M.D., Professor of Materia Medica and Therapeutics, The Thomas W. Evans Museum and Dental Institute School of Dentistry University of Pennsylvania, Philadelphia; formerly Professor of Materia Medica and Therapeutics and Director of the Research Laboratory of the Dental Department of Washington University, St. Louis. 640 pages, 146 illustrations, fourth edition. Price, \$5.00. Published by The C. V. Mosby Company, St. Louis.

This pharmacology constitutes the bulk of his volume and is contained under the heading "Pharmaco-Therapeutics," a term very appropriately coined for the author's intention and accomplishment. That the author himself, lays much claim to this part of his book, appears from the prefaces of his first and subsequent editions.

Dr. Prinz's classification of drugs according to their specific action is a logical sequence of the basal consideration for which we employ drugs at all; namely, their action. For a *materia medica* purely, it is quite immaterial what classification is adopted, but in the pharmacologic sense, where the action of the drugs is primarily concerned, it is but natural that this action should also constitute the basis of their classification. We can scarcely conclude our remarks on this classification without commending the natural and agreeable order in which the author has arranged the various classes. For instance, the antiseptics, covering nearly one hundred pages are given first consideration because they are absolutely of prime importance to the dentist. These antiseptics are immediately followed up by the astringents and the caustics on the principle that the action of these three classes of agents is one chiefly of degree depending on the concentration. The author then introduces successively the chapters on the hemostatics, the demulcents and emollients, the irritants and counterirritants, etc.

While Prinz's "Pharmaco-Therapeutics" constitutes the essence and substance of his volume, yet, this part is not introduced until the author has acquainted us with Part I which he entitles "General Therapeutics" and in which he seizes the opportunity of laying a solid foundation for the better understanding of the succeeding subjects. This part contains the author's introduction which displays his ability as a writer and affords sufficient interest to merit its reading a number of times on account of the historical data contained in it. The chapters on the "Aims of Therapeutics," "Classification of Dental Remedy," "Methods of Administering Medicines," "Prescription Writing," etc., are well and ably written with a view of imparting thorough and reliable information.

These two parts thus briefly reviewed would constitute the subject matter of a *materia medica* proper, but Dr. Prinz's book being, besides, a treatise on dental therapeutics must, necessarily, also consider those remedial agents which do not participate in the characteristic of material substances. These agents are considered under the heading of "Physical-Therapeutics" which forms Part III of the volume and covers some fifty pages. The leading subjects of this part are those on "Artificial Hyperemia," "Massage," "Light Therapy," "Heat and Cold," and a few others. Those subjects are all treated in their most recent conception and acceptance. The author also introduces under this heading of Physical-Therapeutics, the new and very interesting subject on "Ionic Medication." Unfortunately, in the opinion of the reviewer, this subject should not have been included under the general heading of Physical-Therapeutics, because its treatment at this point creates the impression as though ionic medication is essentially a pharmacological one, the employment of the electric current being only a means of transporting the *material* ions into the tissues of the body. The treat-

ment of the subject, therefore, belongs rather to Pharmaco-Therapeutics than to Physical-Therapeutics.

The author has reserved for a separate part (Part IV) of his volume the very important subject of "Local Anesthesia" to which he devotes not less than seventy pages. This part is a complete and thorough exposition of every detail of local anesthesia, both in principle and in technic. Conductive anesthesia, so much talked about in recent times, is also fully considered at this point and we might add here, that this phase of anesthesia has appeared in all former editions of Prinz's, so that it is not a new subject in this book. Dr. Prinz's treatment of local anesthesia may be rightly considered as one of his strong points.

In the new edition, many of the subjects have been completely rewritten while a large number of others have received extensive alterations and additions of new matter, among which we note the following articles: "Constituents of Organic Drugs," "National Narcotic Laws," "Biologic Test for Arsenic," "Formalin Dermatitis," "Formaldehyd as a Desensitizing Agent," "Emetin in the Treatment of Pyorrhea Alveolaris," "Quinin and Urea Hydrochlorid," "Ionic Medication," and "Radio-active Substances."

We cannot conclude our review without taking due notice of the manner in which Dr. Prinz supplements his writings with fitting illustrations. The number of illustrations accompanying the text is one hundred and forty-nine. A crude figure on the black board of a triangle assists the mathematician in his intricate and extended calculations, although the dimensions of the lines are purely imaginary and do not enter into his speculations. So it is with a fitting illustration in a text; it guides the imaginative faculty which directs the mental thought. Personally the reviewer attaches great value to these illustrations, a characteristic feature of Prinz's book. The reviewer is reminded of a Dental Congress from which one seldom departs without taking with him some fresh resolutions which were engendered right in the exhibition hall. Last, and not least, we laud the publishers for the mechanical execution of the new volume which is perfect in every respect.

INDEX TO VOLUME II.

AUTHORS' INDEX

A

- ANGLE, E. H. Some new forms of orthodontic mechanism, 584

B

- BAKER, LAWRENCE W. Orthodontia—Its place in dental education, 148
— Further evidence as to the influence of the forces of occlusion in the development of the bones of the skull, 524
BARR, W. G. The evolution of the treatment of distocclusion, 252
BEASER, HARRY P. A "patent" suggestion, 701
— A removable retaining appliance, 708
BLILER, H. E. Treatment of certain types of malocclusion with the aid of bridge-work, 703
BRADFORD, EDWARD H., and LOVETT, ROBERT W. Orthopedic surgery, 123
BROTHERS, ELMER D. Jurisprudence, 81
BROWN, JOHN MACKENZIE. Report of a case of chronic suppuration of the antrum in which the radiographic examination was indispensable, 470

C

- CASE, CALVAN S. The bodily movement of the teeth in orthodontia, 519
CHAMBERLAIN, W. A. Preventive dentistry, 349
COSTON, W. A. Bands or crowns for temporary molars, 705

D

- DARLING, BYRON. Oral sepsis as a focus of infection, 295
DEWEY, MARTIN. (*See* Skinner and Dewey), 451
— Intramaxillary anchorage, 696
— The lingual arch in combination with the labial arch with extensions as used by Dr. Lloyd S. Lourie, 593
— The principles of anchorage, 539
— The teeth as factors in the economy of the animal kingdom, 138
— The use of the lingual arch in the treatment of malocclusion as used by Dr. Lloyd S. Lourie, 648
DICK, J. LAWSON. The teeth in rickets, 585
DUCKWORTH, T. G. Inheritance of malocclusion from a biological standpoint, 157

E

- EBY, JOSEPH D. The evolution of the principles of appliances, 184, 246, 330, 481
ELLIS, WALTER H. An appliance for opening up space for unerupted bicuspids, 217
— A report of a case treated with the pin and tube appliance, 263
ENDLEMAN, JULIO, and MCCOY, JAMES D. Clinical and radiographic evidence that crown and bridge-work is a menace, 510
ENGSTROM, CARL O. A unique method of mounting photographs, 342

F

- FEDERSPIEL, M. N. A new method for the relief of lateral tension in cleft palate operation, 475
FERNALD, ADELBERT. Students as orthodontists, 69
FISK, EUGENE LYMAN. The role of mouth infections and mouth abnormalities in the causation of disease, 693

G

- GIFFORD, A. C. Compromise treatment, or conservative orthodontia, 197
GUILFORD, S. H. Methods of teaching orthodontia to dental students, 168.

H

- HAMILTON, H. B. A report of an accident, 712
— Changing from general practice to orthodontia, 215
HANAU, RUDOLPH L. Dental engineering, 533
HECKER, F. A plea for conservation of the cementum, 91
— Histological studies of the development of the alveolus of young Rhesus monkeys, 605
HELLMAN, MILO. A further study of some etiological factors of malocclusion, 218.
HOWARD, CLINTON C. The result of unphysiological pressure expressed in two cases, 602

J

- JACKSON, D. E. An experimental investigation of the pharmacological action of nitrous oxide, 544
- JOHNSON, C. N. Textbook of operative dentistry, 122

K

- KELSEY, HARRY E. An effective method for the mesial or distal movement of individual teeth in the arch, 13
- KETCHAM, A. H. Dr. Angle's new pin and tube appliance, 171
- Treatment by the orthodontist supplementing that by the rhinologist, 15
- KEYS, FREDERICK A. (*See* Littig and Keys), 672
- The radiogram as a factor in better dental surgery, 623
- KNOCHE, KARL G. Other points in malocclusion, 121

L

- LEACH, FLOYD D. A symposium on dental radiographic technic, 467
- LEWIS, SAMUEL J. The technic of accurate impression taking, 29
- LISCHER, B. E. Face facts. A clinical study of dento-facial deformities, 129, 187, 397
- LITTIG, MARK D., and KEYS, FREDERICK A. Norman Greene Reoch, D.M.D., 672
- LOEB, VIRGIL. A study of the cubic capacity and superficial area of the maxillary sinus, 639
- LOVETT, ROBERT W. (*See* Bradford and Lovett), 123

M

- MARTIN, W. W. The ideal age for the correction of malocclusion, 212
- MCCARTER, W. A. The development of the mandible as a result of treatment of the protruding upper incisors and distoclusion of the arch, 668
- MCCOY, JAMES DAVID. A consideration of some of the factors involved in the interpretation of dental and oral radiographs, 572
- (*See* Endelman and McCoy), 510
- Suggestions in x-ray technic for the orthodontist, 1
- The interpretation of dental and oral radiographs, 285
- The nature of the x-ray and its discovery, 385
- MCKAY, FREDERICK S. A peculiar accident, 537
- MENDEL, LAFAYETTE B., and OSBORNE, THOMAS B. Growth, 616

- MITCHELL, EDWARD L. Is facial development an index to character? 675

O

- OSBORNE, THOMAS B. (*See* Mendel and Osborne), 616

P

- POLLOCK, H. C. Case history, 118
- The practical use of tungsten and molybdenum in orthodontic appliances, 661

R

- ROSENOW, EDWARD C. Elective localization of the streptococcus from a case of pulpitis, dental neuritis and myositis, 713
- RUCKSTUHL, URLING C. A case of neurotrosis, complicated by extreme distorsion of the upper central incisors, and redundancy of number, 162

S

- SEYDELL, E. M. Infections of the antrum, 344
- SKINNER, E. H., and DEWEY, MARTIN. A comparative study of the attachment of the teeth of lower animals, 451
- STANTON, FREDERICK LESTER. Orthodontic engineering, 235

T

- THOLEN, E. F. Report of a case of dentigerous cyst of the mandible in which the x-ray proved of great value, 517
- Some instances in the practice of oral surgery where the x-ray has been of great value, 580
- THOMPSON, HAROLD B. Plea for a more general use of the x-ray, 726
- TOUSEY, SINCLAIR. Prevention of malocclusion, 680

W

- WEBSTER, RAYMOND L. Gaining and keeping the child's confidence during orthodontic treatment, 165
- WEINBERGER, BERNHARD W. The history of orthodontia, 40, 103, 203, 264, 359, 424, 488, 562
- WEISS, OSCAR A. Dental colleges in relation to the teaching and practice of orthodontia, 315
- WILLETT, R. C. Model library index, 201
- Modern office appointment for the orthodontist, 356, 409
- WOODS, FRANK R. Retention, 486, 535

Z

- ZENTLER, ARTHUR. The factor of heredity in malocclusion, 177

GENERAL INDEX

A

- Abnormal dentition, 192
- Abnormal facial types, 187
- Abnormal function of lips, 190
- Abnormalities of mouth, role of, in causation of disease, 693
- Accident, dental, report of, 712
- Adenoid operation, not always cure for mouth-breathing, 15
- Age, ideal, for correction of malocclusion, 212
- Alignment of teeth, normal variations of, 132
- Alinement and malalignment, casts showing, 193
- Alveolar abscesses, 296
- Alveolus of young monkeys, development of, histological studies of, 605
- American Medical Association, president of, friend of dental profession, 472
- American Medical Association convention, dental exhibit at, 474
- American Society of Orthodontists, members of, 523
 - president's address, 689
 - proceedings of, 586
 - program of, 531
 - sixteenth annual meeting of, 586
- Anchorage, principles of, 539
 - compound primary intramaxillary, 699
 - intramaxillary, 697
 - methods of, 697
 - primary, 699
 - reinforced intramaxillary, 697
 - simple, 697
 - simple primary intramaxillary, 699
 - stationary, 700
- Angle's method for determining profile proportions, 400
- Angle's pin and tube appliance, 171, 584
- Antrum, chronic suppuration of, radiographic examination in, 470
 - infection of, 344
 - of Highmore, 639
- Animal experiments for localization of streptococcus, 715
- Apparatus, dental surveying, 235
- Appliance for opening up space for unerupted bicusps, 217
- Appliances, evolution of principles of, 184, 246, 330, 481
 - fixed and removable, differentiated, 185
 - ideal, requisites of, 184
 - removable, retaining, 708
 - universal regulating, 731
 - use of tungsten and molybdenum in, 661
 - small gauge regulating, 736

- Attachments of teeth of lower animals, comparative study of, 451
- Attitude of dental profession toward orthodontia, 173

B

- Baker anchorage, 256
- Bands versus ligature, 61
- Bands or crowns for temporary molars, 705
- Bilateral distocclusion, defined, 195
- Bilateral mesiocclusion, defined, 195
- Biology and orthodontia, 157
- Blind abscess, 297
- Bodily movement of teeth in orthodontia, 519
- Breast-feeding versus bottle-feeding in their influences upon development of teeth, statistics on, 218
- Bridgework, removable, consideration of problems involved in, 121
 - treatment of malocclusion with aid of, 703
- Buccoversion, defined, 195

C

- Caries, and malocclusion, 321
 - in childhood, 688, 741
- Case history, Class II, Division 1, 118
- Case's method for determining profile proportions, 403
- Cementum, a plea for the conservation of, 91
- Cephalic index, 130
- Changing from general practice to orthodontia—a few efforts, 215
- Child's confidence during orthodontic treatment, gaining and keeping of, 165
- Chin, prominent, with normal denture, 134
 - receding, with normal denture, 135
 - straight, with normal denture, 135
- Chin-cap, devised by Cellier, 265
 - devised by Fox, 266
- Clamp band, first application of, 448
- Class II, Division I, case history of, 118
- Classification of head measurements, 130
 - of malocclusion in Forsythe Dental Infirmary Clinic, 128
 - of malocclusion, Stanton's method, 238
 - of malposition of teeth, 195
- Cleft palate operations, new method for relief of lateral tension in, 475
- Clinical and radiographic evidence that crown and bridge-work is a menace, 510
- Clinical diagnostic aids in orthodontic treatment, 404
- Clinical malocclusion, 126

Clinical orthodontia, 126
 Coffin's split-spring plate, 246
 Columbia University establishes first university dental school in New York City, 176
 Comparative study of attachments of teeth of lower animals, 451
 Compound intramaxillary anchorage, 697
 Compound primary intramaxillary anchorage, 699
 Compromise treatment, or conservative orthodontia, 197
 Confidence of child, gaining and keeping, during orthodontic treatment, 165
 Conservative orthodontia, 197
 Co-operation between dentist and orthodontist, 57
 Correction of malocclusion, ideal age for, 212
 Cretinism and malocclusion, 220
 Crimes of orthodontia, 589
 Crown and bridgework is a menace, clinical and radiographic evidence, 510
 Crowns or bands for temporary molars, 705
 Cubic capacity and superficial area of the maxillary sinuses, 639
 Cyst of the mandible, value of x-ray in, report of a case, 517

D

Dental accident, report of, 712
 Dental caries in childhood, 688
 Dental chart for diagramming composite x-ray findings, 306
 Dental colleges, in relation to the teaching and practice of orthodontia, 315
 teaching of orthodontia in, 63
 Dental disaster, 309
 Dental engineering, 235, 533
 Dental hygienists, new law provides, 622
 Dental laws and the state board, 311
 Dental libraries, 232
 Dental literature—is it free from commercial influences? 734
 Dental profession's attitude toward orthodontia, 173
 Dental societies and ownership of papers presented before them, 687
 Dental surgery, radiograph as factor in better, 623
 Dental surveying apparatus, 235
 Dentist and orthodontist, co-operation between, 59, 315
 Dentist, personality of, 175
 Dentition, abnormal, 192
 normal variations of, 132
 Dento-facial deformities, clinical study of, 129, 187
 Denture, device for outlining, 235, 533
 Development, facial, an index to character, 675

Development of mandible as a result of treatment of protruding upper incisors and distoclusion of the arch, 668
 Diagnosis in dentistry, value of x-ray in, 726
 Distal movement of individual teeth in the arch, an effective method for, 13
 Distoclusion case, report of, 670
 Distoclusion, evolution in treatment of, 252
 Distoclusion of arch and treatment of protruding upper incisors, development of mandible as result of, 668
 Distoversion, defined, 195

E

Eastern Association of Angle Graduates, 7th annual meeting, 284
 Educating the public on orthodontic subjects, 318, 324, 529
 Elective localization of the streptococcus from a case of pulpitis, dental neuritis, and myositis, 713
 Engineering, dental, 533
 orthodontic, 235
 Esthetic side of orthodontia, 65
 Etiological factors of malocclusion, 218
 Evolution of principles of Jackson's appliances, 185, 246, 330, 481
 Evolution of treatment of distoclusion, 252
 Experimental investigation of pharmacological action of nitrous oxide, 544
 Extramaxillary anchorage, 543

F

Face, principal features of, 136
 Face facts: a clinical study of dento-facial deformities, 129, 186, 397
 Face-form and head-form, relation between, 133
 Facial angle, 137
 Facial development an index to character, 675
 Facial index, 134
 Facial orthopedia, 123
 Facial types, abnormal, 187
 Factor of heredity in malocclusion, 177
 Family traits, as factor in inheritance theory of malocclusion, 159
 Farrar's method for determining profile proportions, 399
 Federspiel's tension plates for cleft palate operation, 476
 Finger springs, 338
 Forces of occlusion, influence of, on development of bone of skull, 524
 Forsythe Dental Infirmary, undergraduate assistants for, 211

G

Gnathic index, 131
 Goode, Dr. William, obituary notice of, 737

Growth, 616
 Grünberg applicator, 31
 Grünberg groover, 31

H

Habits, perverted, as cause of malocclusion, 178
 Harvard University Dental School, teaching of orthodontia in, 148
 Head measurements, 130
 Head-form, relation to face-form, 133
 variation of, 130
 Heredity in malocclusion, the factor of, 177
 Heredity of malocclusion from biological standpoint, 157
 Histological studies of development of alveolus of young monkeys, 605
 History of orthodontia, 40, 103, 202, 264, 359, 424, 488, 562
 Hygiene, dental, 251
 Hygiene of head, important field of medicine, 693

I

Ideal age for correction of malocclusion, 212
 Impression taking:
 fitting the trays, 30
 for making retainers, 39
 instruments used in, 30
 lower impression, 36
 modeling compound, not suitable for, 29
 plaster of Paris, best material for, 29
 preparation of patient, 29
 technic of accurate, 29
 to ascertain normal mesio-distal relation of molars, 39
 upper impression, 32
 Watson diagram for, 31, 36
 where teeth are missing, 38
 Index, model library, for orthodontist's records, 201
 Infections of the antrum, 344
 Infections of the mouth, role of, in causation of disease, 693
 Infraversion, defined, 195
 Inheritance of malocclusion from a biological standpoint, 157
 Instruments, sterilization of dental, 120
 Inter-marriage of races as factor in producing malocclusion, 159
 Intermaxillary anchorage, 543
 Internal secretions, important etiological factor of malocclusion, 230
 Interpretation of dental and oral radiographs, 285, 572
 Intramaxillary anchorage, 696
 Invention in dentistry, 701

J

Jackson's removable appliances, 185, 246, 330, 481

Journal of National Dental Association, value of, to dental profession, 631
 Journals, state dental, value of, 473
 "Jumping the bite," 256
 Jurisprudence, dental, 91

L

Labioversion, defined, 195
 Lane groover, 31
 Lane separator, 31
 Langenbeck's method for relief of lateral tension in cleft palate operations, 475
 Lateral tension, new method for relief of, in cleft palate operations, 475
 Library index for orthodontist's records, 201
 Life extension institute, dental inspection in, 696
 Ligature versus bands, 61
 Lingual arch in combination with labial arch with extensions as used by Dr. Lloyd S. Lourie, 593, 648
 Linguoversion, defined, 195
 Lips, abnormal functions of, 190
 malformation of, 191
 malrelation of lower to upper, 190
 Literature, dental, and commercial influences, 734
 Loop appliance, question of priority in use of, 175

M

Macrogathia, defined, 196
 Malalignment, defined, 192
 Malformations of the jaw and their processes, defined, 192
 Malocclusion:
 classification, in Forsyth Dental Infirmary Clinic, 128
 effect of, on the beauty of the face, 66
 etiological factor of, 218
 hereditary, 157
 and occlusion, cast showing, 193
 factor of heredity in, 177
 perverted habits, 178
 ideal age for correction of, 212
 in dental clinics, 126
 in school children, 159
 inheritance of, from biological standpoint, 157
 treatment of, with aid of bridgework, 703
 Malocclusions and cretinism, 220
 Malposition of teeth, defined, 192
 forms of, 195
 Malposition of the mandible, defined, 192
 Malrelation of the dental arches, defined, 192
 Mandible, development of, as result of treatment of protruding upper incisors and distocclusion of arch, 668

- Materia medica and therapeutics for dentists, 738
- Maxillary sinuses, study of cubical capacity and superficial area of, 639
- Measurement, x-ray, of unerupted permanent teeth to provide for regulation of temporary arch, 680
- Median maxillary suture, opening of, 22
- Mesial movement of individual teeth in the arch, an effective method for, 13
- Mesioversion, defined, 795
- Micrognathia, defined, 196
- Mixing of types, as factors in producing malocclusion, 159
- Model library index for orthodontist's records, 201
- Models, accurate value of, 29
 - unique method of photographing, 670
- Modern office appointment for the orthodontist, 356, 409
- Molars, lower third, variations in relative positions of, 132
 - normal mesio-distal relation of, impression to ascertain, 39
- Molars, temporary, bands or crowns for, 705
- Molybdenum, in orthodontic appliances, practical use of, 661
- Mounting photographs, unique method of, 342
- Mouth-breathing, adenoid operation not always cure of, 15
 - prevalence of, 15
- Mouth hygiene, importance of, 693
- Mouth infection and abnormalities, role of in causation of disease, 693
- Movement of individual teeth an effective method for, 13
- Movement of teeth in orthodontia bodily, 519

N

- Nasal variations affecting the profile, 398
- National Dental Association, Journal of, value of, to dental profession, 631
 - research institute of, 176
- Nature of x-ray and its discovery, 385
- Neuroclulsion, case of, complicated by extreme distoversion of the central incisors, and redundancy of number, 162
- Nitrous oxide anesthesia, 547
 - action of, 552
 - symptoms produced by, 549
- Nitrous oxide apparatus, 545
- Nitrous oxide, experimental investigation of pharmacological action of, 544
- Nonocclusion, 193, 195

O

- Occlusion and malocclusion, cast showing buccal aspect, 193

- Occlusion of permanent teeth of man, 194
- Office appointment for the orthodontist, 356, 409
- Operating room illumination, 329
- Oral sepsis as a focus of infection, 295
- Oral surgery, relation to orthodontia, 124
 - some instances in practice of, where the x-ray has been of great value, 580
- Orthodontia:
 - and general physical health, 294
 - and post-graduate instruction, 391
 - and process patents, 394
 - its place in dental education, 148
 - prosthetic dentistry, and crown and bridgework, how related to, 231
 - study of, a most interesting and instructive branch of dentistry, 68
- Orthodontic engineering, 235, 533
- Orthopedic surgery and orthodontia, 123

P

- Parathyroid, disturbances of, effect on teeth, 222
- Pathology, dental, and roentgenography, 684
- Partial clasps, construction, 332
- Patent board for National Dental Association suggested, 702
- Pathology, dental, x-ray in diagnosis of, 726
- "Patent" suggestion, 701
- Patients, education of, 529
 - ethical phase of referring, in orthodontic practice, 526
- Pediatrician and orthodontist, 634
- Periodental membrane, 91
- Periosteum, 91
- Permanent teeth, unerupted, x-ray measurements of, to provide for regulation of temporary arch, 680
- Personality of dentist, 175
- Perversion, defined, 195
- Perverted habits, cause of malocclusion, 178
- Photographing models, unique method of, 670
- Photographs, unique method of mounting, 342
- Pin and tube appliance, Angle's, 171
 - report of case treated with, 263
- Pituitary body, disturbances of, effect on the teeth, 224
- Plaster of Paris, as impression material, 29
- Plea for more general use of x-ray in dentistry, 726
- Post-graduate instruction and orthodontia, 391
- Prenatal causes of malocclusion, 178
- Preventive dentistry—an answer to the present dental dilemmas, 349
- Primary anchorage, 542, 699
- Principles of anchorage, 539

- Principles of appliances, evolution of, 184
- Prinz, new book on materia medica and therapeutics, 738
- Priority in use of loop appliance, 174
- Process patents and orthodontia, 394
- Profile, evolution of, 130
 - methods for determining proportions of, 398
 - nasal variations affecting, 398
- Prosthetic dentistry, orthodontia, and crown and bridge work, how related, 231
- Prosthodontia, specimens of work from prominent men, 302
- Protruding upper incisors and distoclusion of the arch, development of the mandible as result of, 668
- Pyorrhea alveolaris, conservation of cementum in, 91

R

- Radiograph, as factor in better dental surgery, 623
 - of cyst of mandible, 517
- Radiographic diagnosis of dental infection in systemic diseases, 628
- Radiographic technic, symposium on, 467
- Radiographs, dental and oral, interpretation of, 285, 572
 - examination of, 287
- Radiography (*see* X-ray)
- Radiography, terminology, 685
- Raymond's method for determining profile proportions, 398
- Reciprocal intramaxillary anchorage, 543
- Referring patients in practice of orthodontia, 526
- Regulating appliances, 731, 736
- Reinforced intramaxillary anchorage, 697
- Removable appliances, evolution of, 184, 246, 330, 481
- Removable retaining appliances, 708
- Reoch, Dr. Norman Greene, obituary notice of, 672
- Report of case treated with pin and tube appliance, 263
- Report of a dental accident, 712
- Research Institute of National Dental Association, 176
- Retainers, impressions for making, 39
- Retaining appliance, removable, 708
- Retention, 486, 535
 - intermaxillary, 256
- Rhinologic and orthodontic treatment, 15
- Rhinologist and orthodontist, relationship between, 528
- Rhinologist and orthodontist should work together, 26
- Rickets, teeth in, 585
- Röntgen ray in dental practice, 630

- Röntgenography and dental pathology, 684
- Röntgenologist, mechanic or consultant, 637
- Roots of teeth, localized infections of, 713
- Role of mouth infections and mouth abnormalities in causation of disease, 693

S

- Scientific subjects before orthodontic societies, a plea for more careful consideration of, 588
- Sexual glands, disturbances of, effect on teeth, 225
- Small gauge regulating appliances, 736
- Simple anchorage, 542
- Simple primary intramaxillary anchorage, 699
- Sinuses, maxillary, cubic capacity and superficial area of, 639
- Skull, influence of forces of occlusion in development of bones of, 524
 - measurement of, 130
- Societies, dental, and ownership of papers presented before them, 687
- Spring clasps, 333
- Spring force for tooth movement, 246
- St. Louis dental society, 395
- Stanton's classification of malocclusion, 238
- State boards and dental laws, 311
- State dental journals, 473
- Stationary anchorage, 700
- Stationary intramaxillary anchorage, 543
- Sterilization of dental instruments, 120
- Streptococcus, localization of, from a case of pulpitis, dental neuritis, and myositis, 713
- Streptococcus, fermentative powers of, 724
- Students as orthodontists, 69
- Supraversion, defined, 195
- Suture, median maxillary, 22
- Surveying apparatus, Stanton's, 235
- Symposium on dental radiographic technic, 467

T

- Taggart inlay case, appellate court decision in, 591
- Teaching and practice of orthodontia, dental colleges in relation to, 315
- Teaching of orthodontia in dental colleges, 63, 168
- Teeth and tuberculosis, 310
- Teeth as factors in the economy of animal kingdom, 138
- Temporary molars, bands or crowns for, 705
- Terminology in radiographic work, 685
- Thymus, disturbances of, effect on teeth, 223

- Thyroid gland, disturbance of, effect on teeth, 222
 Tonsils, benefit to, from orthodontic work, 26
 Torsoversion, defined, 195
 Transversion, defined, 195
 Trays in impression taking, fitting of, on patient, 30
 Treatment by orthodontist supplementing that by rhinologist, 15
 Treatment, method for estimating in advance probable effect of, 405
 Tungsten and molybdenum, in orthodontic appliances, practical use, 661

U

- Unilateral distoclusion, defined, 195
 Unilateral mesioclusion, defined, 195
 Universal regulating appliance, 731
 Unphysiological pressure, result of, expressed in two cases, 602

V

- Value of orthodontic treatment, 529

W

- Watson diagram for impression taking, 31, 36
 Wiegall's method for determining profile proportions, 398


X

- X-ray, nature and discovery of, 385
 X-ray in dentistry:
 arrangement of apparatus, 2
 dental chart for diagramming findings, 306
 extra-oral method, 5
 intra-oral method, 5
 photographic dark room, 3
 plea for more general use of, 726
 position of patient, 9
 technic for orthodontists, 1, 3
 terminology, 685
 value of, 1, 285, 304
 X-ray measurement of unerupted permanent teeth at the age of five or six years to provide for regulation of temporary arch if required, 680
 X-ray plates, as court evidence, 471
 examination of, 287


EDITORIALS

- A new fourth edition of Prinz's "Dental Materia Medica and Therapeutics," 738
 A plea for more careful consideration of scientific subjects before orthodontic societies, 588
 American Institute of Dental Teachers, 638
 "Apropos," 68
 Bands vs. ligature, 61
 Co-operation between the dentist and the orthodontist, 59

- Dental caries in childhood, 688
 Dental laws and state board, 311
 Dental libraries, 232
 Dental societies and the ownership of papers presented before them, 687
 Dr. William Goode, 737
 First university dental school in New York for Columbia, 176
 Is our dental literature free from commercial influence? 734
 Malocclusion in dental clinics, 126
 National Dental Association's action endorsed, 234
 Orthodontia and post-graduate instruction, 391
 Orthodontia and process patents, 394
 Permanent staff appointments for the Forsyth Dental Infirmary for Children, 233
 Preventive orthodontia, 633
 Prosthetic dentistry, orthodontia, and crown and bridge work, 231
 Referring patients, 526
 Relation of oral surgery to orthodontia, 124
 Report on classification in the Forsyth Dental Infirmary Clinic, 128
 Sixteenth annual meeting of the American Society of Orthodontists, 530
 Sixtieth anniversary meeting of the St. Louis Dental Society, 592
 State dental journals, 473
 The Appellate Court decisions of the Taggart inlay case, 591
 The attitude of the dental profession toward orthodontia, 173
 The crimes of orthodontia, 589
 The dental exhibition at the American Medical Association Convention, 474
 The department of dental and oral radiography, 313
 The esthetic side of orthodontia, 65
 The Journal of the National Dental Association, 631
 The new president of the American Medical Association, 472
 The personality of the dentist, 175
 The president's address before the fifteenth annual meeting of the American Society of Orthodontists, 689
 The proceedings of the American Society of Orthodontists, 586
 The question of priority, 174
 "The roentgenologist: mechanic or consultant," 637
 The St. Louis Dental Society, 395
 The small gauge regulating appliance, 736
 The teaching of orthodontia in dental colleges, 63
 The universal regulating appliance, 731
 The value of orthodontic treatment, 529



*The Problem of
Tooth Rotation Solved
with*



Angle's New Ribbon-Arch Orthodontic Appliances



[Patented]

Rotation has always been the most difficult of all tooth movements because complete force-control was impossible with the various forms of mechanism formerly employed.

Dr. Angle's New Ribbon-Arch Appliances permit every form of root and crown movement—rotation, depression, elevation as well as mesial and distal movement in the line of occlusion.

These appliances are easily manipulated, positive in action, cause no tissue disturbance, are light and comfortable—meet every physiological and hygienic need.

Angle's New Ribbon-Arch Orthodontic Appliances are a revelation in regulating, require less frequent appointments, obtain results satisfactory to patient and operator.



Write for reprint in pamphlet form of Dr. Angle's article in the September 1916 issue of the "Dental Cosmos," which describes these appliances in detail.

Sent with price list, free upon request.


For Sale by Dental Dealers and at our Houses

The S. S. White Dental Manufacturing Co.

"Since 1844 the Standard"

Philadelphia

New York, Chicago, Boston, Atlanta, Brooklyn, San Francisco, Oakland





COLUMBIA



FIFTY-FIVE MODERN DENTAL OFFICES

is the title of a book we have just completed, showing floor plans and other details of fifty-five selected offices, we have prepared for the profession.

Our experience in this work will enable us to be of assistance to you in solving problems of this kind, by drafting detailed plans and offering suggestions on the matter of furnishings and decoration to fit your particular case.

A copy of this book of plans containing as well a detailed explanation of this service will be sent with our compliments upon receipt of request and dealer's name.

THE RITTER DENTAL MFG. CO.
ROCHESTER, N.Y.


CHICAGO

PHILADELPHIA

NEW YORK

B-114

Does the Dentist need the X-Ray?



Even yet this question is sometimes asked! The answer is: all the progressive dentists—all those who are making a reputation—are putting in X-ray Equipment.

The Fischer Outfit is a simple practical and efficient X-Ray machine and equipment, designed on *modern* lines, expressly for Dental use.

No rectifiers; no acid interrupters; no valve tubes; no inverse. Not a single moving part—so nothing to wear out.

Thoroughly endorsed and guaranteed.

Sold by the best
Dental Dealers
everywhere.

Send for full descriptive circular
and get name of
nearest Agency.



H. G. Fischer & Co. 2343 Wabansia Ave.
CHICAGO

The Right X-RAY Apparatus! *What should you buy?*



If you want to make dental radiographs only you can get a suitable equipment equal to any for quality of work at less than \$250.00.

If you intend to do frontal sinus and other difficult work and contemplate buying an Interrupterless Machine, be sure that machine has the necessary spark gap to use a tube of such penetration. The best work will be done with a tube backing up a seven inch spark. Forget about the toys with a five to six inch spark gap. Buy a real machine.

If you are interested in either equipment write us.

THE WM. MEYER COMPANY

Washington Boulevard and Green Street
DEPT. O. - - - CHICAGO, ILL.

ORTHODONTIC APPLIANCES AND SUPPLIES DENTAL SPECIALTIES

**A postal card brings our illustrated
catalogues to your office.**


**A new and enlarged edition of our
catalogue is in preparation.**

**Our Goods are Sold by Dental Dealers
Everywhere**

BLUE ISLAND SPECIALTY CO.
MANUFACTURERS

BLUE ISLAND, ILL., U. S. A.

IMPRESSION TRAYS



Complete set of Upper and Lower Impression Trays, in handsome box, either mahogany or ebony finish. Price, \$15.00.

PRICE OF SINGLE IMPRESSION TRAYS

Nos. 1, 2, 3, 4, 5, 6, 18, 19, 20, 21, 22, 23.....	\$0.50
Nos. 7, 8, 9, 10, 11, 12, 24, 25, 26, 27, 28, 29.....	.40
Nos. 13, 14, 15, 16, 17, 30, 31, 32, 33, 34.....	.35

Blue Island Specialty Co., BLUE ISLAND, ILL.
U. S. A.

¶ Treat Pyorrhea by the Dunlop Method

¶ Effective and Speedy Cure
the Result

¶ You Should Investigate
This Form of Treatment


¶ *Descriptive Literature outlining
method of treatment sent upon
request*

¶ *Address*

WILLIAM F. DUNLOP,
M.D., D.D.S.

24 East 48th Street, New York City, N. Y.

New Revised Fourth Edition of Prinz Now Ready



DENTAL MATERIA MEDICA AND THERAPEUTICS

With Special Reference to the Rational Application of Remedial Measures to Dental Diseases.—A Text Book for Students and Practitioners.

By HERMANN PRINZ, A.M., D.D.S., M.D.

Professor of Materia Medica and Therapeutics, The Thomas W. Evans Museum and Dental Institute School of Dentistry University of Pennsylvania; formerly Professor of Materia Medica, Therapeutics, and Pathology, and Director of the Research Laboratory, Washington University Dental School, St. Louis.

Octavo, 634 pages, with 149 engravings.—Fourth revised edition—Cloth, \$4.00

THE demand for a fourth edition of "Dental Materia Medica and Therapeutics" has necessitated a complete revision of the text. The "Introduction" and the chapters on The Classification of Dental Remedies, Decomposition of the Tooth Pulp and Its Treatment, Preparations for the Mouth and Teeth (Oral Hygiene), Uric Acid Solvents and Local Anesthesia have been completely rewritten, while many other chapters have received extensive alterations. The chapter containing articles on Constituents of Organic Drugs, National Narcotic Law, Formaldehyd as a Desensitizing Agent, Formalin Dermatitis, Biologic Test for Arsenic, Emetin in the Treatment of Pyorrhea Alveolaris, Quinin and Urea Hydrochlorid, Ionic Medication and Radio-Active Substances have been newly added, while the chapter on Urine Analysis has been omitted. The present edition is increased by some fifty odd pages with the addition of some fifty new illustrations.

It is seldom that critics and reviewers accord such unusual praise to a book as has been given to Dr. Prinz' Dental Materia Medica and Therapeutics. This laudation is international in extent, and is due to the high character and scientific quality of the matter contained in this book. The matter is mostly original, and the result of a quarter of a century of effort in the laboratory, in private practice, and in university teaching is embodied in it. The tribute that has been paid it by teachers and critics at home and abroad is the credit due to a work well done. The book is the most valuable contribution to the science of dentistry that has been made in a decade.

Send your order to your dealer or direct to publisher.

The C. V. MOSBY COMPANY, Dental Publishers
METROPOLITAN BUILDING, Grand Avenue and Olive Street ST. LOUIS, U. S. A.



A clean mouth is an essential of good health. In the attainment of a clean mouth Dioxogen deserves consideration.

Combining definite germ destroying properties with detergent qualities of a high order, Dioxogen is a product that every dentist should be acquainted with.

The Oakland Chemical Co.

10 Astor Place

NEW YORK, N. Y.

DENTAL X-RAY FILMS

PARAGON X-RAY PLATES

WE CALL YOUR ATTENTION to the fact that we are ready to supply you, from stock carried in Chicago, or other large cities, or through any reputable dealer,

PARAGON X-RAY PLATES. Used by leading Orthodontists everywhere. Noted for extreme speed, combined with a rich black quality, showing wonderful detail, and enabling the X-Ray worker to make a quick and sure diagnosis.

DENTAL X-RAY FILMS carried in either fast or slow emulsion, in all standard sizes, including ovals. Some special sizes, such as 4x5 and 5x6 inches.

FILM DEVELOPING HOOKS, three kinds. **DENTAL FILM MOUNTS**, in either celluloid or paper. **DEVELOPER** in tin cans, ready for use by dissolving in water. **ACID FIXING BATH.**

SEND US YOUR ORDERS and write us about your X-Ray troubles. We maintain the **ONLY X-RAY INFORMATION BUREAU.** Ask for our free book on X-Ray. If you send a plate order, we will furnish a marker with your name, **GRATIS**, on request.



GEO. W. BRADY & CO.

776 S. WESTERN AVE.
CHICAGO, ILL.



Model 1D



DENTAL X-RAY APPARATUS
The Coil of Quality

CONVENIENT SAFE SIMPLE

Combines

X-Ray - Violet Ray

Switchboard Instruments

**THE PRICE IS RIGHT
THE TERMS ARE
RIGHT**

**Detach and Mail
This Coupon
To-Day**

Gentlemen—Please send literature. I am
interested in
Dental X-Ray Transformer.
Name.....
Address.....

Campbell Elec. Co., Lynn, Mass.

A New Book on an Important Subject


DENTAL ANATOMY

By

MARTIN DEWEY, D.D.S., M.D.

PROFESSOR OF DENTAL ANATOMY AND ORTHODONTIA, KANSAS CITY
DENTAL COLLEGE; PRESIDENT OF THE DEWEY SCHOOL
OF ORTHODONTIA, KANSAS CITY, MO.

300 pages, with 133 New and Original Engravings
Price, silk cloth binding, \$2.50




SPECIMEN ILLUSTRATION

The C. V. Mosby Company—Publishers
St. Louis, U. S. A.

Edwards Perfected Dental X-Ray Apparatus

Machine, Tube Stand and Protective Screen complete in one unit. When not in use the tube carrying arm is swung above, and the screen folded flat against the cabinet. The floor space required is no more than for cabinet alone. This is only one of the many advantages of the Edwards machine.

You
CAN
PAY
MORE
but you
cannot
BUY
A
BETTER
machine



Thorough investigation will prove TO YOU that this machine HAS NO EQUAL in efficiency, simplicity and durability AT ANY PRICE.

Designed and constructed by pioneers in the X-Ray industry.

GUARANTEED
FOR FIVE YEARS

Our latest catalogue, illustrating Dental X-Ray machines from \$125.00 up, is just off the press. Write for your copy to-day. Please mention your dealer's name and this Journal.

Edwards Instrument Company
968 Ft. Wayne Avenue INDIANAPOLIS, IND.

You Need Not Inflict Pain in Your Work
Use
SOMNOFORM
For
ANALGESIA OR ANÆSTHESIA



It will solve the problem for you. Pain can be eliminated. Better, quicker, more thorough work can be done. All this has been proven by SOMNOFORM. Our lessons will help you to quickly master the technique.

Write for free text book

Send me your free Text Book, on SOMNOFORM. Enclosed find 60 cents for which send lessons No. 1, 2, 3 and 4 of your Post Graduate Course.

Name.....

Address

Dealer's Name.....

STRATFORD-COOKSON CO.

Successors to

E. De TREY & SONS,

28 South 40th Street

PHILADELPHIA . . . PA.

"LIFE IS A CONSTANT STRUGGLE
AGAINST OXYGEN DEFICIENCY"
(Pawlow)

CALOX

The Oxygen Tooth Powder

(McK&R)

is more than a dentifrice, more than a potential germicide or antiseptic,
more than a pleasant toilet necessity.

CALOX liberates *Nascent Oxygen* which destroys pathogenic organisms
as well as products of fermenting or putrefying organic matter.

Therefore, CALOX is a Prophylactic agent which appeals to dentist,
doctor, surgeon or specialist, who are all, or should be all, alive to the
part oral sepsis plays in the production of general severe or even fatal
disease.


Samples on request.

McKESSON & ROBBINS

ESTABLISHED 1833

NEW YORK

Made in the (McK&R) laboratories.



McIntosh Dental Ionization Switchboard

McIntosh Battery & Optical Co.,
217-223 N. Desplaines Street, CHICAGO, ILL.

Dental Ionization

Every progressive dentist nowadays is
talking ionization. Are you prepared to
apply this method in your work? If not,
you need this switchboard. Let us send you
a description of same and a special
price good during December only.

FREE! An interesting re-
print on Ionization. Use
the coupon or mail a
post card.

Mc-
Intosh
Battery &
Optical Co.,
Chicago, Ill.

Gentlemen:
You may send me
free reprint on Ioniza-
tion and description and
price of your Switchboard.

Name.....

Address.....

I. J. of O.

THE VULCAN COIL



Type L

This beautiful Colonial apparatus, so designed as to show, when closed, no hint of its purpose may be obtained in three sizes, 2 KW, 3 KW and 4 KW. power, the largest occupying a floor space of only twenty by twenty-four inches, with all the Simplicity, Reliability and EFFICIENCY of the original Vulcan Coil, which has made many a Radiographer his reputation.

MATCHLESS

In power and speed under twice the price, Guaranteed FIVE YEARS—will last a lifetime. Fast and powerful enough for any purpose in radiography—no subject beyond its range—the smallest machine producing high class intraoral radiographs in four seconds, the largest in ONE SECOND.

SEND FOR CATALOG

Catalog showing various models, with full description and sample radiographs mailed on request, together with address of nearest agency.

Vulcan Coil Company

241 South Los Angeles St.

Los Angeles, California

Vulcan Coil Co., 241 S. L. A. St., Los Angeles, Cal.
Name
Address

CONTENTS

Original Articles

- The Role of Mouth Infections and Mouth Abnormalities in the Causation of Disease. By *Eugene Lyman Fisk, M.D.*, New York 693
- Intramaxillary Anchorage. By *Martin Dewey, D.D.S., M.D.*, Kansas City, Mo. 696
- A "Patent" Suggestion. By *Harry P. Beaser, D.D.S.*, Fresno, Calif. 701
- Treatment of Certain Types of Malocclusion With the Aid of Bridge-Work. By *H. E. Bliler, D.D.S.*, Chicago, Ill. 703
- Bands or Crowns for Temporary Molars. By *W. A. Coston, D.D.S.*, Topeka, Kansas. 705
- A Removable Retaining Appliance. By *Harry P. Beaser, D.D.S.*, Fresno, Calif. 708
- A Report of an Accident. By *H. B. Hamilton, D.D.S.*, Ithaca, N. Y. 712

Department of Dental and Oral Radiography

- Elective Localization of the Streptococcus from a Case of Pulpitis, Dental Neuritis and Myositis. By *Edward C. Rosenow, M.D.*, Rochester, Minn. 713
- Plea for a More General Use of the X-Ray. By *Dr. Harold B. Thompson*, Seattle, Wash. 726

Editorials

- The Universal Regulating Appliance. 731
- Is Our Dental Literature Free From Commercial Influence? 734
- The Small Gauge Regulating Appliance. 736
- Dr. William Goode. 737
- A New Fourth Edition of Prinz's "Dental Materia Medica and Therapeutics" 738

1917 Announcement

The
DEWEY SCHOOL
of
ORTHODONTIA

is a highly specialized institution whose course is so arranged as to promote orthodontia as a science and a specialty. The curriculum is such as to be thoroughly adequate to meet the requirements of those wishing to practice orthodontia as a specialty or teach the same in colleges. At the same time, for the man expecting to include orthodontia in his general practice, thorough equipment of the operator is the first essential, and we believe a compromise course has no place in modern scientific orthodontia.

The next session will be held July 2nd, 1917, and continue for eight weeks. Applications for the 1917 session should be made early in order to insure a place, as the class is limited.

*Prospectus of the school and further information
may be obtained direct from*

H. C. POLLOCK, D.D.S., Secretary

808 Metropolitan Building,

St. Louis, Mo.

or

MARTIN DEWEY, M.D., D.D.S.

1016 Armour Boulevard

Kansas City, Mo.

LAVORIS

A PERFECTLY REPAIRED OROPHYLACTIC REPARATION **A PARTICULARLY REFERRED PROFESSIONAL DESCRIPTION**

FOR

MOUTH-THROAT-NOSE

FIRST AID TO THE DENTIST

LAVORIS CHEMICAL COMPANY
MINNEAPOLIS, MINN., U.S.A.

RADIOGRAPH "PHILMOUNTS"


Increase the Detail of Film (Picture)
A Protection to Emulsion Side of Film
A Means of Filing and Recording

(Trade Mark)
Eliminate Foreign Objects when Viewing
A convenient Holder when Viewing
An Advertisement when Sent Out

CASE NO. _____

M _____ **DATE** _____

(VIEW ONLY BY STRONG TRANSMITTED LIGHT.)



LINGUAL ASPECT

JAMES T. CASE, M. D.
Roentgenologist

Roentgen Department
BATTLE CREEK SANITARIUM

PATENT PENDING, SWENARTON STATIONERY COMPANY, NEW YORK

Send for Sample and Price List 14 Stock Sizes Special Sizes to Order
SWENARTON STATIONERY COMPANY, 121 East 27th St., New York, N. Y.
Publishers of F. LeRoy Satterlee's "Dental Radiology." \$3.00, postpaid.

HEIDBRINK AUTOMATIC ANAESTHETIZER

By *Accurate Control* of gases and *Markings* on *dials* indicating dosage for anaesthesia and analgesia

NITROUS OXIDE-OXYGEN

anaesthesia is greatly simplified.

*Write for further
information.*

THE HEIDBRINK CO.
Minneapolis, Minn.

When you clean teeth, do it by TAXI

The word TAXI is copyrighted for your protection. The naming of the world's greatest tooth cleaning preparation is a happy one, for it conveys more nearly than any other word in the dictionary would, exactly the popularity of TAXI. You find TAXI everywhere. TAXI is universally used and TAXI does the work for you. It serves. Every bottle of TAXI has TAXI Service in it.

TAXI cleans the teeth of all gangrene, calcareous deposits, tobacco stains, removes deposits of all kinds.

TAXI is used before examination of teeth for small cavities which are often obscured by deposits. TAXI is used after dental operations leaving the teeth sweet and clean.

TAXI is the first aid to dentists who know about it and last resort for those who fail without it.

Use TAXI—it is strictly up-to-date.

Full directions on every bottle.

Price \$1.00 per bottle—6 ounces

For Sale By All Progressive Dealers

The Sam'l A. Crocker Co.

Established 1872

Incorporated 1911

Makers of the celebrated Crocker Rolling Mills. Publishers of the Dental Register


OHIO DENTAL AND SURGICAL DEPOT

Conrad Building, 18-20 West Seventh Street

CINCINNATI, OHIO

In the Heart of the Dental Zone

EXODONTIA



The New Book
For the Exodontist and
General Dental
Practitioner

BY

GEORGE B. WINTER, D.D.S.,

Professor of Exodontia and Lecturer on
Anesthesia, St. Louis University
School of Dentistry

Large Octavo Size - - 430 Pages
251 Original Illustrations

Price - - - \$4.00

THE BEST WORK ON EXTRACTION EVER PUBLISHED

It is often necessary to extract teeth. Sometimes this becomes a difficult and dangerous operation. It is at such times as this that you will need a book that tells you how to master such difficulties and obviate the dangers. In Dr. Winter's book you get graphic detailed description of just how to meet these cases and master them. Never before have such beautiful and instructive illustrations been incorporated in a book on this subject.

If you do not possess a copy of this book order it today

CONTENTS

- | | |
|---|--|
| I—Historical. | X—Extraction Technic of the Inferior Teeth. |
| II—Instruments. | XI—Extraction Technic of the Inferior Third Molar. |
| III—Office Equipment. | XII—Deciduous and Supernumerary Teeth. |
| IV—Anatomical Landmarks. | XIII—Hypercementosis and Artificial Complications. |
| V—Indications and Counter-Indications for Extraction. | XIV—Accidents. |
| VI—Examination of the Mouth. | XV—Treatment after Extraction. |
| VII—Position of the Patient and Operator. | XVI—Hemorrhage. |
| VIII—Precautionary Suggestions. | XVII—General and Local Anesthesia. |
| IX—Extraction Technic of the Superior Teeth. | |

Sent prepaid, on receipt of price

ST. LOUIS MEDICAL BOOK COMPANY

508 North Grand Avenue

ST. LOUIS, MISSOURI

Meet Me at the **TULLER**
For Value, Service, Home Comforts



New HOTEL TULLER DETROIT, MICHIGAN

Center of business on Grand Circus Park. Take Woodward car, get off at Adams Avenue

ABSOLUTELY FIREPROOF

200 Rooms, Private Bath, \$1.50 Single, \$2.50 up Double
200 " " " 2.00 " 3.00 " "
100 " " " 2.50 " 4.00 " "
100 " " " 3.00 to 5.00 " 4.50 " "

Total, 600 Outside Rooms—All Absolutely Quiet
Two Floors—Agents' Sample Rooms
New Unique Cafes and Cabaret Excellente

Hotel Muehlebach

BALTIMORE AVENUE AND TWELFTH STREET
Kansas City, Mo.



500

New Fireproof Rooms


Rate from \$2.00

The House of
Utility—Service—Elegance

Operated By
Whitmore Hotel Co.
Under the Personal Direction of
S.J. Whitmore and Joseph Reichl

Jackson's Orthodontic Appliances

A SPECIALTY



Made for the Dental Profession by an expert with seven years' experience in Dr. Jackson's office. There have been many improvements in the last two years and every day my mail increases from dentists wishing to give the Jackson System a trial. Are you one of them?

Price, \$7.00

GET THE ORIGINAL AND BE SAFE

CHESTER T. SWEET

SPECIALIST IN JACKSON REGULATING APPLIANCES

17 West 42nd Street

Telephone, 2278 Vanderbilt

NEW YORK CITY

THE LARYNGOSCOPE

An International Monthly Journal
Devoted Exclusively to Diseases of the

EAR—NOSE—THROAT

Subscription Price, \$5.00 per Annum, in Advance

To recognize the various conditions in their relation to diseases of the ear, nose or throat the orthodontist should keep thoroughly posted on the literature of these specialties. In addition to important original communications, society proceedings and abstracts each month, *The Laryngoscope* publishes annually a complete Index Medicus of the literature of Otology, Rhinology and Laryngology, with abstracts of the most valuable articles in these specialties.

Make subscription remittance payable to

THE LARYNGOSCOPE CO.

ST. LOUIS, MO.

3858 WESTMINSTER PLACE

E. J. FAGAN

7 West 45th Street - - New York

REFINER OF

GOLD AND SILVER

MANUFACTURER OF

Dental Gold Plate and Solders

Send me your Old Gold and Grindings. We will refine and return either a Check, Gold Plate, Solder or Shells

Try Me

Agent for Daly's Gold Lining

SALESMEN WANTED

We now have openings for several salesmen to call on Physicians and Dentists. Exclusive territory. New and attractive line. Our representatives make money.

C. V. MOSBY COMPANY

MEDICAL PUBLISHERS

Metropolitan Bldg.

ST. LOUIS

SAL HEPATICA

Materially
AIDS

Local Treatment
In
PYORRHEA

Bristol-Myers Co.
New York





SMITH'S COPPER CEMENT is not sold on a competitive basis as to price.

It is being used exclusively by dentists throughout the world *solely* on account of its superior quality.

But because it is **UNIVERSAL** and performs *all* the functions of *all* the oxyphosphate of zinc cements and *all* the functions of *all* the copper cements, it is supplied in a

FULL ASSORTMENT PACKAGE

containing six each double portions of powder and liquid, the colors assorted any way you wish.

Price of the full assortment package is \$15.00.

The same amount of material purchased in single color packages, a box at a time, would cost you \$24.00.

And when you use *one* cement for *all* classes of work you have but *one* technique to master.


But **SMITH'S COPPER CEMENT** is the *only* cement that is **UNIVERSAL** and that may be used for *all* classes of work because it is the *only one* that contains more than enough copper to render it a perfect germicide, yet

WILL NOT DISCOLOR IN THE MOUTH

Made in 8 shades in the laboratories of

LEE S. SMITH & SON MFG. CO.,
PITTSBURGH, U. S. A.

Recommended in Leading Dental Colleges



PRACTICAL DENTAL METALLURGY

A Text and Reference Book for Students and Practitioners
of Dentistry

Embodying the Principles of Metallurgy and
Their Application to Dentistry,
Including Experiments

FOURTH EDITION—COMPLETELY REVISED

By JOSEPH DUPUY HODGEN, D.D.S., Professor of Operative
Dentistry (Formerly Professor of Dental Chemistry and Metallurgy),
Dental Department University of California. Revised by GUY
S. MILLBERRY, D.D.S., Professor of Chemistry and Metallurgy,
Dental Department University of California.

374 Octavo Pages, 44 Illustrations, PRICE \$2.50

Table of Contents

- I. Introduction.
- II. The Properties of Metals.
- III. Combination of Metals with Non-Metallic Elements.
- IV. Melting Metals.
- V. Alloys.
- VI. Lead.
- VII. Antimony.
- VIII. Tin.
- IX. Bismuth.
- X. Zinc.
- XI. Cadmium.
- XII. Copper.
- XIII. Iron.
- XIV. Aluminum.
- XV. Mercury.
- XVI. Silver.
- XVII. Iridium.
- XVIII. Palladium.
- XIX. Platinum.
- XX. Gold.
- XXI. Amalgams.
- XXII. Classified Amalgams.

Recommended by Leading Colleges

The matter contained in this instructive text-book, which is now in its fourth edition, is the result of the author's work in the laboratory and his experience as an instructor on the subject of Dental Metallurgy. The book presents in a clear and practical manner the principles of the subject as the author sees them related and applicable to the every-day wants of the dentist. The subject is so treated that the book may be taken into the Metallurgical laboratory and used as a manual for practical and experimental work.

This is the fourth edition of this book, and has been thoroughly revised, presenting the newest facts and most accurate data obtainable, thus keeping the work abreast with the scientific investigations of the day.

It is safe to say that Hodgen's Dental Metallurgy is the text-book which is best adapted to the needs of the student and practitioner. This fact has been recognized by the majority of dental colleges, who, in consequence, have placed the work upon their list of required text-books.—Dental Era.

THE C. V. MOSBY CO., Dental Publishers
801-807 Metropolitan Building SAINT LOUIS, U. S. A.

CRESCENT ALLOY

FOR PERFECT FILLINGS

The New Gould's Practitioner's Medical Dictionary

3d Edition

Entirely Reset from New Type

70,900 Terms

Includes Words in Medicine
and Allied Sciences

Clear, Readable Type

Contains the Current Words
and Terms

Pronounced and Defined
On the Famous Gould System
Accurate and Reliable
Definitions Concise and Clear
Gives Derivation

Tables and Summaries for Quick
Reference 962 Pages, Weighing
only 2½ Pounds

Flexible Bindings, Cloth and
Leather, Marbled Edges,
Round Corners

Height, 8¾ Inches; Width, 6¼
Inches; Thickness, 1¼ Inches

Handsome Flexible Cloth, Mar-
bled Edges, Round Corners, \$2.75,
or with Thumb Index, \$3.25.

Beautiful Flexible Leather, Mar-
bled Edges, Round Corners, \$4.00,
or with Thumb Index, \$4.50,
postpaid.

.....CUT HERE.....

ON APPROVAL ORDER BLANK

P. BLAKISTON'S SON & CO.

Date _____

1012 Walnut St., Philadelphia.

Please send me postpaid, for examination, a copy
of Gould's Practitioner's Dictionary, 3d Revised
Edition, 70,900 words, Cloth \$2.75 (Indexed \$3.25)
Leather \$4.00 (Indexed \$4.50.)

Name _____

Address _____

Mark Style Desired

I. J. O.

California Dental Quiz and Post Graduate Course

Preparatory to State Dental Examination
of Pacific Coast States including Arizona
and Nevada. Army and Navy Examina-
tions: Civil Service for City, County and
State Institutional and School Work.

*The Post Graduate Feature is Alone Worth
the Price of Tuition*

NEXT QUIZ BEGINS APRIL 26, 1917

A. HALDEN JONES, A.M., M.D.
222 Bradbury Bldg., Los Angeles, Cal.

The Sexual Life

By C. W. MALCHOW, M.D.

Formerly Professor of Proctology and
Associate in Clinical Medicine, Hamline
University College of Physicians and
Surgeons; President of Physicians' and
Surgeons' Club. 318 pp. Third edition.
A Scientific Treatise Designed for Ad-
vanced Students and the Professions,
Embracing the Natural Sexual Impulse,
Normal Sexual Habits, and Propagation,
Together with Sexual Physiology and
Hygiene. Postpaid, on receipt of price,
\$3.00.

In this book we find a very satisfactory dis-
cussion of the sex problem in all its phases,
in masterly and chaste English, and every
one who reads it will profit by it. We have
often been amused by physicians asserting that
"everybody knows all about this subject." If
you think you know all about the sex prob-
lems, buy this book and you will discover
some new light on an old subject.—*The Med-
ical Herald.*

The C. V. Mosby Co.
Publishers SAINT LOUIS

Bound Volumes

WE can still supply a number of bound
volumes of Vol. I, 1915, issue of
International Journal of Orthodontia at
\$5.00 per volume in black silk cloth,
with name stamped in gilt letters on back-
bone of volume.

The C. V. Mosby Co.
Publishers SAINT LOUIS

CHLORAZENE

Dakin's New Antiseptic

(Para-toluene-sodium-sulphochloramide)

This interesting antiseptic was introduced by Dr. H. D. Dakin, who was associated with Dr. Alexis Carrel at the temporary hospital No. 21 and the laboratories of the Rockefeller Foundation, in Compiègne, France. Reports show this antiseptic is being used with great success in the military hospitals of Europe. (See British Medical Journal of January 29, 1916. Reprints on request.)

CHLORAZENE promises to be of great value in surgical practice. Read the following statements and it will be understood why this new antiseptic is regarded so favorably in Europe.

It is a white crystalline substance, freely soluble in water.

It is extremely stable, i.e., in solid form it may be preserved indefinitely, while aqueous solutions keep for a relatively long time without marked decomposition.

It has no corrosive action, even in concentrated solutions. It neither precipitates nor coagulates proteins such as blood serum.

It is virtually non-toxic. Rabbits and guinea-pigs tolerate subcutaneous doses of a Gram to a kilo (2 1-5 pounds) of body-weight, with no symptoms except moderate local reaction. However, it should not be used internally.

Its antiseptic action is intense. It is many times as antiseptic as phenol, equalling sodium hypochlorite in this respect (molecule for molecule it is four times as germicidal as hypochlorite), while, according to Dakin, "it is much less irritating than the latter substance, and may be used safely at a concentration five to ten times as great."

It can be used freely and safely as an irrigant of infected wounds of deep tissues, in treating compound fractures and in injuries and diseases of the mouth, uterus, bladder and urethra, where most antiseptics cannot be employed in effective concentration on account of their toxic or caustic action.

In view of the success which has followed the use of this antiseptic in Europe—

The Abbott Laboratories offer it to the medical profession of America under the name **CHLORAZENE** (Abbott).

Literature on Request.

PACKAGES AND PRICES

CHLORAZENE is supplied in 4.6-grain tablets, in bottles of 100 at 60c. In powder; two special packages for general and hospital use: Hospital Package No. 1, to make 1 gallon of 1-per cent solution, 55c; Hospital Package No. 2, to make 5 gallons of 1-per cent solution, \$2.00. Prices on larger quantities on request.

The trade will be stocked, but if your druggist is not supplied, we shall be glad to supply you direct, from our home office or branches.

THE ABBOTT LABORATORIES

CHICAGO - NEW YORK

EATTLE

SAN FRANCISCO

LOS ANGELES

TORONTO

BOMBAY

THE WINTER SCHOOL OF EXODONTIA

The Scientific Extraction of Teeth
CONDUCTED BY GEORGE B. WINTER, D. D. S.
Exodontia Specialist

AN EXCEPTIONAL OPPORTUNITY

The Winter School of Exodontia presents an exceptional opportunity for dental graduates, whether practitioners or teachers, to perfect themselves in Exodontia, the scientific extraction of teeth.

INDIVIDUAL PRIVATE INSTRUCTION

There will be no class instruction in this School, as each person taking the course will be given individual private instruction in Exodontia, including the administration of nitrous oxid and oxygen, the technic of general and local anesthesia, and practical radiography.

The instruction will be by the personal demonstration of Dr. Winter, and will be conducted in such a systematic and practical manner as to properly qualify the individual as an Exodontist.

For further Information Address:

GEORGE B. WINTER, D.D.S.

906 OLIVE STREET

ST. LOUIS, MO., U. S. A.

BEFORE and AFTER

DENTURE is inserted in mouth; no place to fill up with soft tissue and become irritated as in the old worn-out methods.



BEFORE INSERTING



AFTER INSERTING

Thousands of Dentures are in daily use with the "EUREKA" attached that could never wear a plate before.

CAN YOU AFFORD TO TAKE CHANCES?

PRICE—\$2.00 PER BOX—HALF DOZEN

EUREKA SUCTION CO.
LOUDONVILLE, OHIO

Sherman's Bacterial Vaccines

Preparations With A Record For
EFFICIENCY

In the treatment of mouth infections, Bacterial Vaccines are now recognized as most efficient remedies. This will include mandibular, septic apical and dento-alveolar abscesses, and pyorrhea. Sherman's List No. 40 or 35 are admirably adapted for these uses.

Sherman's New All Glass Syringe

An Unusual Value, Price \$2.00

Two Platin Needles, Self-Sterilizing Cap over Needle and Metal Case



Patent Applied for

Write for Literature

G. H. SHERMAN, M. D. DETROIT,
U. S. A.

AUTOGENOUS VACCINES FOR PYORRHEA AND BONE INFECTIONS

Wassermann Tests

ALLEN H. BUNCE, M.D.,

824 HEALY BUILDING

ATLANTA, GA.

A Book Every Dental Practitioner Should Have

SUGGESTIVE THERAPEUTICS Applied Hypnotism and Psychic Science


By HENRY S. MUNRO, M.D.
Omaha, Neb.

New Fourth Edition, Revised and Enlarged.
Original Frontispiece

410 Royal Octavo Pages—Price, \$5.00

Sent Anywhere, Express Prepaid, on Receipt of Price

Doctor Munro has written the most practical book on Suggestion that has been published. That this book is of great value to the dentist is evidenced by the large sale it has had among the dental profession. The author has very successfully stated facts in a simple way, which are ordinarily couched in technical language, and has added many illustrative cases to render things still more clear. He deserves great credit for his work, and we believe every dental practitioner, who is not already familiar with the cardinal principles of psychotherapy, will gain much by a perusal of this work. Sound, common sense is the keynote of the whole work. This edition has been thoroughly revised, eight new chapters have been added, and new typography has been used throughout. It is eminently the helpmeet of the general practitioner, and has placed within his reach one of the most valuable therapeutic adjuncts.




The C. V. MOSBY COMPANY, Medical Publishers

METROPOLITAN BUILDING, Grand Avenue and Olive Street

ST. LOUIS, U. S. A.

NEW POINTS *for the* HARPER HOLDER

To meet the growing demand I present a set of Enamel Cleavers Nos. 1, 2, 3, and Root Trimmers Nos. 4, 5, 6, 7, designed by



DR. FORREST H. ORTON, of St. Paul, Minn.,

for use in the Dental Department of the University of Minnesota.

The reduced size and smoothly rounded form of the working point and the additional angle of shank gives to this set of seven points distinct merit, and greatly increases the range and possibilities of perfect root preparation for the dentist who strives for PERFECTION in his crown and bridgework.

PRICES

Harper Holder and the Cleavers, Nos. 9, 10, 11.....	\$3.00
Orton Cleavers and Trimmers Nos. 1, 2, 3, 4, 5, 6, 7, each, 50c; set.....	3.50
Cleavers: No. 9; Right No. 10; Left, No. 11, Each.....	.35

May be purchased through your dealer or from

DR. WM. E. HARPER

Manufacturer of

Harper's Quick Setting Alloy. The Harper Holder and Cleaver Points.
The Universal Trimmer, and Inventor of the Contra Angle Hand Piece.

3441 Michigan Avenue,

CHICAGO, ILL.

Orthodontic Appliances

Dental Laboratories

*Full and Complete Line of
Dental Supplies*

Hettinger Bros. Mfg. Co.

Kansas City, Mo.

Oklahoma City, Okla.

St. Louis Dental Mfg. Co.

(Hettinger Bros.)

St. Louis, Mo.

Why Tire Yourself Out?

The Noratchet Operating Stool

Automatically
Meets Every
Motion



Saves Nerves
Preserves Health
Increases
Working Time
When Slightly
Inclined
Chest is Thrown
Forward and
Shoulders Back

A pleasing, agreeable and esthetic, as well as a most sensible and practical addition to the equipment of any office, having no projecting levers; its mechanical features are such as skill, ingenuity and long experience suggest, leaving nothing to be desired.

While the seat revolves freely, such revolution has nothing to do with its height, that feature being controlled by a gripping device within the standard which cannot slip under the weight of the operator. The seat is adjusted up or down by a touch, the adjusting lever being a simple steel ring under the seat, entirely out of the way, yet always just where either hand drops naturally, wherever that may be. No pushing is required, since there are no springs to overcome. It is transported about the office by tipping slightly and rolling upon its base.

The Noratchet Operating Stool Will Help

PRICES

Noratchet Operating Stool.....	\$20.00
Stock Finishes—Decorated black, gray, mahogany, blue and white enamel.	
*Stock Upholstery—Maroon and green plush; green blue, gray and maroon hair-cloth; green, black, maroon, gray and red leather.	
Special finishes and upholstery, other than above, extra.....	\$3.00
Special finish only, other than above, extra.....	2.00
Special upholstery only, other than above, extra.....	1.00
Extra height for men over six feet tall, extra.....	1.50

*Subject to market conditions.

Your Dealer will be glad to Demonstrate THIS Stool and fill your order

The Ransom & Randolph Co.
CLEVELAND TOLEDO GRAND RAPIDS
COLUMBUS

ANNALI DI ODONTOLOGIA

An International Monthly Magazine of Dental Science

edited by ANGELO CHIAVARO of Rome (Italy) with contributions by the most eminent Dentists of the world.

The ANNALI DI ODONTOLOGIA publishes at the end of each original communication and editorial a short summary in Italian, French, English and Spanish, so that the articles can be read and understood by dentists of any nationality.

All the Italian original communications on Dentistry, which appear in other Italian journals, will be published every month summarized in the same four languages.

The eminent contributors, who collaborate to the "ANNALI DI ODONTOLOGIA" are:

Florestan Aguilar, Madrid; Oscar Amoëdo, Paris; R. Anema, Paris; Luigi Arnone, Florence; Aspelund Axel, Helsingfors; Matti Ayrapää, Helsingfors; Arture Beretta, Bologna; Ed. A. Bogue, Chicago; Truman W. Brophy, Chicago; James Cameron, Philadelphia; M. Chivaki, Tokio; C. Christensen, Copenhagen; Lodovico Coulliaux, Milan; Isaac Davenport, Paris; William Davenport, Paris; Beniamino De Vecchis, Naples; M. Dewey, Kansas City; Edward Dunn, Florence; William Dunn, Florence; G. Fiordelmondo, Roma; Ch. Godon, Paris; J. E. Grevers, Utrecht; Vincenzo Guerini, Naples; S. H. Guilford, Philadelphia; P. Guye, Geneva; G. B. Hayes, Paris; Otto E. Inglis, Philadelphia; N. S. Jenkins, New Haven, Conn.; C. N. Johnson, Chicago; Edward C. Kirk, Philadelphia; W. G. Law; G. Lind, Amsterdam; P. Martinier, Paris; Joseph Mendel; J. Howard Mummery, London; Camillo Palazzo, Rome; W. B. Paterson, London; S. B. Piergili, Rome; M. L. Rhein, New York; Maurice Roy, Paris; E. S. Talbot, Chicago; George Villian, Paris; Henri Villian, Paris; H. L. Wheeler, New York; W. J. Younger, Paris.

E. RAMPICHINI, Publisher,—"Annali di Odontologia"

VIA VENTI SETTEMBRE 8, ROMA, (Italy)

Subscription Price, \$1.80 per annum, payable in advance

Send your subscription to the C. V. Mosby Co., Metropolitan Bldg., St. Louis

CAMPHO-PHENIQUE

LIQUID

Invaluable in Dental Surgery



Campho-Phenique (the liquid form) is to be found at the hand of the best dentists in this country. Its uses in the dental office are many, and its worth as a therapeutic agent is highly appreciated by practitioners who are familiar with its virtues.

It is of exceptional value as a local anaesthetic preceding all operations in the mouth. It prevents nausea, obtunds local sensation in removing calculus, extracting loose roots; wedging, and ligating the rubber dam prior to taking impressions, also for treating sensitive cavities preparatory to filling or crowning. It effectually prevents gagging and relieves pain. It is antiseptic, non-irritating, non-poisonous and does not discolor the teeth.

Price, Regular Size \$1.00. Small Size 25c.

Samples mailed free on request



Address **CAMPHO-PHENIQUE COMPANY**
St. Louis, Missouri.

To Those Contemplating Orthodontic Procedure:

YOU no doubt have heard of correspondence schools in other branches of science and trades. You also know the "follow-up" system of correspondence as applied to commercial business.

Did you ever hear of an orthodontist making a business of constructing appliances, who will follow up your case from the beginning to the finish by a thorough system of filed letters, drawings and photographs of your models, a place where you can ask advice, and get help on your case as it proceeds.

Write and send your models to The Ortho. Special Laboratory of Milwaukee, Wis. It costs you nothing for estimates. There is a difference which can not be explained on paper. Perhaps some of your fellow practitioners have been there. Ask them.

Our correspondence and "follow-up" feature adds to the success of your case, and the instruction so gained stimulates you to attempt all kinds of difficult cases.

We instruct you in the minute, detail technic, of adjusting and caring for the appliance.

We keep tab on your doings, and also on the progress of the case, by an ORIGINAL METHOD OF "FOLLOW-UP SYSTEM".

We advise and consult with you throughout the progress, to the finish of the case.

If you have yet to begin your first case in this branch of dentistry we can help you start right.

If you have had but little experience, we would like to add our experience to yours.

If you are a full fledged specialist in this branch we would like to show you what we can do for you in the way of mechanical construction. This means much in this line of appliances.

Reflect And Then Write Us.

The Ortho. Special Laboratory

F. H. BERRY, Manager.

121 WISCONSIN STREET

MILWAUKEE, WIS.

**When visiting New York
make your home at**

Hotel Chelsea

West 23rd Street, at Seventh Avenue

***One of New York's largest, safest and
most refined hotels***

Absolutely Fire proof—European Plan

500 Rooms

400 Baths

Room with adjoining bath \$1.00 and \$1.50

Room with private bath - 2.00

Suites parlor, bedroom and bath
and upward 3.00

Club Breakfast - - - 25c. up

Special Luncheon - - - 50c.

Table d'Hote Dinner - - - 75c.

Cafe Attached

To Reach Hotel Chelsea: From Pennsylvania Station—7th Ave. car south to 23rd St.;
Grand Central—4th Ave. car south to 23rd St.
Lackawana, Erie, Reading, Baltimore and Ohio, Jersey Central and Lehigh Valley R.R. Stations
—take 23rd St. crosstown car east to Hotel Chelsea;

Principal Steamship Piers, Foot West 23rd Street—take 23rd St. crosstown car.

WRITE FOR COLORED MAP OF NEW YORK

Indiana Dental College

An institution that has been teaching the science
and art of Dentistry successfully for over
thirty years

Offers a Three-Year Course

Unexcelled by any dental teaching institution
in the world

Leading to the Degree of D.D.S.

For Catalogue and Booklet address the College at

INDIANAPOLIS, - - INDIANA

Chicago College of Dental Surgery

DENTAL DEPARTMENT VALPARAISO UNIVERSITY

Is the result of over A QUARTER OF
A CENTURY of steady development

The Next Annual Course of Instruction
will begin the First Tuesday in October

Located in the center of the greatest College and Hospital district in the world, with
unlimited clinical facilities. It offers every needed opportunity to the Dental Student.

For Catalog giving details, address

CHICAGO COLLEGE OF DENTAL SURGERY

1747 W. Harrison St.

Chicago, Ill.

Completely Revised—Rewritten—Enlarged

ESSENTIALS OF OPERATIVE DENTISTRY

By W. CLYDE DAVIS, B.S., M.D., D.D.S.

*Dean and Professor of Operative Dentistry and Technic, Lincoln Dental College,
Associated with the University of Nebraska, Lincoln, Nebraska.*

350 Pages—190 Illustrations—Price, \$4.00

Second Revised Edition

CONTENTS:

- | | |
|--|--|
| I. Instrument Nomenclature. | XXI. Manipulation of Cohesive Gold in the Making of Fillings by Classes. |
| II. Cavity Nomenclature. | XXII. Finishing Gold Fillings. |
| III. Cavity Preparation. | XXIII. Manipulation of Amalgam in the Making of a Filling. |
| IV. Gaining Access. | XXIV. The Use of Cements in Filling Teeth. |
| V. Outline Form. | XXV. Manipulation of Synthetic Porcelain in the Making of a Filling. |
| VI. Resistance Form. | XXVI. The Use of Gutta-Percha in Filling Teeth. |
| VII. Retention Form. | XXVII. Tin as a Filling Material. |
| VIII. Convenience Form. | XXVIII. Combination Fillings. |
| IX. (A) Removal of Remaining Carious Dentine. | XXIX. Examination of the Mouth Looking to Dental Services. |
| (B) Finishing Enamel Walls. | XXX. The Alleviation of Dental Pains. |
| (C) Toilet of the Cavity. | XXXI. Prophylactic Treatment of the Mouth. |
| X. Management of Pit and Fissure Cavities. Class One. | XXXII. Exclusion of Moisture. |
| XI. Management of Pit and Fissure Cavities. Class One Included. Large Cavities in Central Fossa of Molars. | XXXIII. The Treatment of Hypersensitive Dentine. |
| XII. Management of Proximal Cavities in Bicuspids and Molars. Class Two. | XXXIV. Protection of the Vital Pulp. |
| XIII. Large Proximal Cavities Endangering the Pulp. | XXXV. Pulp Devitalization and Removal. |
| XIV. Management of Proximal Cavities in Incisors and Cuspids not Involving the Angle. Class Three. | XXXVI. Management of Putrescent Pulp Canals. |
| XV. Management of Proximal Cavities in Incisors Involving the Angle. Class Four. | XXXVII. The Filling of Pulp Canals. |
| XVI. Management of Cavities in the Gingival Third. Class Five. | XXXVIII. Management of Children's Teeth. |
| XVII. Management of Abraded Surfaces. Occlusal and Incisal. Class Six. | XXXIX. Extraction of Permanent Teeth. |
| XVIII. Cavity Preparation for Gold Inlays. | XL. Extraction of Temporary Teeth. |
| XIX. The Making and Setting of a Gold Inlay. | XLI. Local Anesthesia. |
| XX. Manipulation of Cohesive Gold in the Making of a Filling. | XLII. The Use of Fused Porcelain in Filling Teeth. |
| | XLIII. Preparation of Cavities for Porcelain Inlays. |
| | XLIV. The Construction and Placing of a Porcelain Inlay. |

THE C. V. MOSBY CO. Publishers
801-807 Metropolitan Building - - - ST. LOUIS

Dental Radiographic Laboratory

Special Equipment for
Orthodontic Work
Unusual care taken
to produce results
of the highest
order of merit

CONSULTATION
BY APPOINTMENT

Gray Chandler Briggs, M.D.

WALL BUILDING
SAINT LOUIS, MO.

Lindell 5600

PHONES

Delmar 4206

Orthodontic and Dental Radiographic Laboratory

Special equipment
for rapid and care-
ful dental work

A. E. BELTON

Metropolitan Building
ST. LOUIS

Lindell 5470

PHONES

Central 4175

AUTOTOXEMIA

Is unquestionably one of the most constant causes of dental caries and the early loss of the teeth. Intestinal elimination is, therefore, a fundamental detail of any effective treatment—or prophylaxis—and in selecting measures for this purpose, the dentist will find

PRUNOIDS

(EDIBLE TABLETS)

the solution of a most important problem. Happily the administration of this ideal product is not merely attended by catharsis. Its effects are much more far reaching, and used occasionally PRUNOIDS will restore the functional activity of the whole intestinal canal. The glandular structures are stimulated, the muscles are toned and adequate elimination assured.

The remarkable freedom of PRUNOIDS from griping or reactionary constipation, and its physiological stimulation of intestinal processes, make it the most satisfactory laxative that the dentist can employ.

Liberal samples on request.

THE SULTAN DRUG CO.

ST. LOUIS, MO.

Sold by Druggists.

The Last Word on Radiography

Dental and Oral RADIOGRAPHY

By JAMES DAVID McCOY, D.D.S.

*Professor of Orthodontia and Radiography, College of Dentistry, University
of Southern California, Los Angeles, California*

157 Pages — 71 Illustrations — Price, \$2.00

CONTENTS

- | | |
|--|---|
| I. The Nature of the X-Ray and Its Discovery. | IV. Requisites of the Dental X-Ray Laboratory. |
| II. High Tension Electric Currents—Magnetism — Electro-Magnetic Induction. | V. The Technic of Dental and Oral Radiography. |
| III. X-Ray Machines. The Rhumkroff or Induction Coil—The Tesla or High Frequency Coil—The Interrupterless Transformer. | VI. The Technic of Dental and Oral Radiography. |
| | VII. The Interpretation of Radiographs. |
| | VIII. The Dangers of the X-Ray and Methods of Protection. |

Read What the Reviewers Say

It is always a distinct pleasure to read a book written by an author who knows his business, and who also knows how to tell others about it in a lucid and instructive manner. Röntgenology, like many other ologies, if it is to be studied and comprehended in all of its multitudinous ramifications, relations, and applications, presents today an accumulation of data almost staggering in magnitude. It is a subject beset with obscurities and complexities in its scientific aspects, and with no little danger in its applications when these latter are undertaken by the untrained or ignorant.

On the other hand, there are certain fundamental considerations essential to the intelligent use of this important diagnostic means, which, when clearly presented and grasped, are sufficient to enable those who have mastered them to make intelligent and reasonably safe use of it.

Radiography, as the author very truly states in his preface, was a few years ago considered in the light of a cultural asset to dentistry, but today the farseeing members of our profession have awakened to the fact that it is a real necessity. Dr. McCoy's book "has been written primarily as a textbook for students of dentistry. It is essentially a book for beginners." Of all the attempts which have come under our notice to instruct the novice in the elementary principles and practice of radiography, none in our opinion has so satisfactorily accomplished its purpose as the little volume here under consideration. The electro-physics underlying the phenomena of the x-ray, the means for its production and utilization, the methods for its successful application to the needs of dental practice, are most satisfactorily set forth in a logical and lucid way, so that the student who has mastered the very readable text, even if he have no more than average intelligence, must necessarily know how to safely and successfully utilize this indispensable agency in his practice, and cannot fail to understand the scientific explanations of the phenomenon itself, and the reasoning underlying the technic of its application.—*The Dental Cosmos*.

Send your order to your dental dealer
or direct to the publisher

THE C. V. MOSBY CO., Publishers
801-807 Metropolitan Building - - - ST. LOUIS

Reducing the Price of Teeth AND Not Cheapening the Product IS A Difficult Problem to Solve

We are the first to satisfactorily solve this difficult problem and have placed before the Dental Profession, the well known line of JUSTI TEETH with

"P. M." (Baked in Precious Metal) Pins

A pin that stands every test required of Platinum.

A pin that is Insoluble in Nitric, Hydrochloric and Sulphuric Acids.

A pin that is Higher in tensile strength than one made of Platinum.

A pin that positively does not contain any base metal.

A pin that will save you from 15 to 25% on your tooth bills.

Justi "P.M." Facings

with medium short pins, will (at the present price) save you 25% on Platinum pin facings.

Justi "P. M." Combination Sets

will save you nearly 15% on Platinum pin sets.

Your Dealer can supply you, and will, if you insist on getting "P.M." teeth.

H. D. JUSTI & SON

PHILADELPHIA

CHICAGO

Dentistry's Most Stubborn Problem

SOLVED BY THE USE OF

"PREPAREDNESS"
See your dentist twice yearly
Use Senreco twice daily

The tooth paste that **REALLY CLEANS**

Senreco
Reg. U. S. Pat. Off.



YOU know the problem of saving the anchor teeth that confronts you every time you treat a case of orthodontia.

You fear, in fact are morally certain, that the accumulation of tartar, food, etc., behind the appliances, the difficulty the patient finds in keeping the oral cavity clean, will very likely cause infection and trouble for all concerned.

That problem long confronted one of the foremost men in the profession, one who specializes in Orthodontia and Anaesthetics, one who occupies a chair in one of the foremost dental colleges, who is the author of several recognized works on Orthodontia, and who has invented a number of Orthodontia appliances that are widely used. Here is what he says under date of July 19, 1916:

"I have been prescribing Senreco Tooth Paste for orthodontia patients and I find that it is very efficient in keeping the appliances clean and the gums and contiguous tissues in a healthy condition.

"It is a known fact that keeping orthodontia appliances clean is a difficult task for patients to perform, but I find that those of my patients who have been using Senreco have remarkably clean mouths."

Very truly yours,

(Name furnished upon request.)

Why not put Senreco to that sort of test yourself? We are perfectly willing to have you sit as both judge and jury on the merit of our preparation.

Large size tube and list of ingredients mailed you on receipt of your card.

SENRECO

MASONIC TEMPLE, CINCINNATI, OHIO

The
Post Graduate School of Orthodontia
of the
Forsyth Dental Infirmary for Children

offers an advanced course of thirty-two weeks
from October to May, eclectic in character.

Unexcelled Clinical opportunities.

Broad and thorough theoretical instruction.

A full staff of lecturers.

Correspondence solicited.

Address, NO. 140 THE FENWAY, BOSTON, MASS

ATTENTION TO DENTISTS

It will pay you to send us your Gold Bullion for refining.
Most of the bridge scrap contains some platinum for which the
U. S. Assay Offices are not permitted to make allowance.

We give returns for the full platinum, gold and silver con-
tents and our refining charges are moderate.

We are Refiners and Manufacturers of Platinum, Gold
and Silver and could also furnish you with platinum especially
soft, platinum inlay foil, platinum iridium sheet or wire, either
square or round as well as other shapes. Special alloys of
Platinum, Iridium, Palladium, Gold, etc., made to order in
accordance with specifications.

AMERICAN PLATINUM WORKS

225-231 New Jersey Railroad Avenue, Newark, N. J.

New York Office: CHARLES ENGELHARD, 30 Church Street

SILVOL

A Powerful Non-Toxic, Non-Irritating Germicide, of Marked Value in Dental Practice.

A proteid-silver compound, containing approximately 20 per cent. of metallic silver. Used in the treatment of infections of mucous membranes. Freely soluble in water, and yielding solutions without sediment. Does not coagulate albumin; is not precipitated by proteids or alkalies. Used in solutions of 5 to 25 per cent. without pain or irritation.

♦ ♦ ♦

SILVOL is advantageously used in infection and irritation frequently associated with the correction of irregular teeth; in swabbing cavities following extraction; in painting the gums surrounding decayed teeth. It is applicable, in short, in any irritation or infection of the mucous membranes of the mouth.

One prominent dental surgeon uses SILVOL in 5-per-cent. solution for painting around portions of teeth exposed by receding of the gums. He says: "It works like magic."

SILVOL is practically tasteless.

POWDER: Bottles of one ounce.

CAPSULES (6-grain): Contents of one capsule make one-fourth ounce of a 5-per-cent. solution.

Give SILVOL a trial: its action will not disappoint you. Your druggist or dental depot can supply you.

Home Offices and Laboratories,
Detroit, Michigan.

PARKE, DAVIS & CO.

Washington University Dental School (Missouri Dental College)

The Fiftieth Annual Session will begin September 27, 1915, and continue thirty-two weeks, exclusive of holidays.

This institution offers a thorough course of dental instruction. The advantages offered by a large university like Washington University should appeal to those intending to enter the profession of dentistry. Its location in the fourth largest city in the United States offers unusual clinical advantages.

If you are interested in securing the best there is to be had, do not fail to investigate this institution. For the annual announcement and other information regarding the school, address:

J. H. KENNERLY, Dean,

St. Louis, Mo.

29th and Locust Sts.

